arXiv:hep-ph/0412088v2 30 Mar 2005

Laurent series expansion of a class of massive scalar one-loop integrals to $\mathcal{O}(\varepsilon^2)$.

J.G. Körner[a,†], Z. Merebashvili[b,‡] and M. Rogal[a,*]

a. Institut für Physik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany

b. High Energy Physics Institute, Tbilisi State University, 0186 Tbilisi, Georgia

ABSTRACT

We use dimensional regularization to calculate the $\mathcal{O}(\varepsilon^2)$ expansion of all scalar one-loop one-, two-, three- and four-point integrals that are needed in the calculation of hadronic heavy quark production. The Laurent series up to $\mathcal{O}(\varepsilon^2)$ is needed as input to that part of the NNLO corrections to heavy flavor production at hadron colliders where the one-loop integrals appear in the loop-by-loop contributions. The four-point integrals are the most complicated. The $\mathcal{O}(\varepsilon^2)$ expansion of the three- and four-point integrals contains in general polylogarithms up to Li_4 and functions related to multiple polylogarithms of maximal weight and depth four.

PACS number(s): 12.38.Bx, 13.85.-t, 13.85.Fb, 14.65.Ha

[†] e-mail address: koerner@thep.physik.uni-mainz.de

[‡] e-mail address: zaza@thep.physik.uni-mainz.de

[*] e-mail address: rogal@thep.physik.uni-mainz.de

I. INTRODUCTION

The full next-to-leading order (NLO) corrections to the hadroproduction of heavy flavors have been completed in the late eighties [1,2]. They have raised the leading order (LO) estimates [3] but several initial analysis' showed a serious disagreement with experimental results [4, 5]. Recently the situation has considerably improved in that a more refined NLO analysis (due to considerably more precise experimental input for the b-quark fragmentation function as well as other QCD parameters) now shows signs of rapprochement between theory and the new experimental data (see [6] and references therein for the new CDF measurements). However, the NLO predictions are still slightly below the experimental numbers. Moreover, the theoretical NLO predictions suffer from the usual large uncertainty resulting from the freedom in the choice of renormalization and factorization scales of perturbative QCD. In this light there are hopes that a next-to-next-to-leading order (NNLO) calculation will bring theoretical predictions even closer to the experimental data. Also, the dependence on the factorization and renormalization scales of the physical process is expected to be greatly reduced at NNLO. This would reduce the theoretical uncertainty and therefore make the comparison between theory and experiment much more significant.

In Fig. 1 we show one generic diagram each for the four classes of gluon-induced contributions that need to be calculated for the NNLO corrections to hadroproduction of heavy flavors. They involve the two-loop contribution (1a), the loop-by-loop contribution (1b), the one-loop gluon emission contribution (1c) and, finally, the two gluon emission contribution (1d). A similar classification holds for the quark-induced contributions.



FIG. 1. Exemplary gluon fusion diagrams for the NNLO calculation of heavy hadron production.

In this paper we concentrate on the loop-by-loop contribution Fig. 1b. Specifically, working in the framework of the dimensional regularization scheme [7], we shall present $\mathcal{O}(\varepsilon^2)$ results on all scalar one-loop one-, two-, three- and four-point integrals that are needed in the calculation of hadronic heavy flavour production. We generate the coefficients of the ε-expansion in a rather direct way. Let us briefly describe our procedure. We introduce Feynman parameter representations for each of the two-, three- and four-point integrals. The two-point case is straightforward. In case of the three- and four-point integrals we generally integrate over one and two Feynman parameters, respectively, keeping the full ε-dependence of the result. Before doing the last Feynman parameter integration we expand the respective integrands up to $\mathcal{O}(\varepsilon^2)$ and then integrate the expanded integrand term by term.

Because the one-loop integrals exhibit ultraviolet (UV) and infrared (IR)/collinear (or mass (M)) singularities up to $\mathcal{O}(\varepsilon^{-2})$ one needs to know the one-loop integrals up to $\mathcal{O}(\varepsilon^2)$ because the one-loop contributions appear in product form in the loop-by-loop contributions[1]. It is clear that the spin algebra and the Passarino-Veltman decomposition of tensor integrals in the one-loop contributions also have to be done up to $\mathcal{O}(\varepsilon^2)$. This task will be left to a companion paper where we present complete results on the one-loop amplitudes up to $\mathcal{O}(\varepsilon^2)$ including spin algebra and Passarino-Veltman decomposition effects.

The general case of massive one-loop integrals was studied some time ago [9], where a general one-loop N-point integral was expressed in terms of hypergeometric functions of several variables. Recently, there have been a number of papers where the authors took a more general attitude to calculate the ε-expansion of massive one-loop integrals. They write down general representations of the ε-expansion of one-loop integrals for general kinematic configurations. We have attempted to compare our results with the results of the more general approaches whenever possible. In papers [10, 11] the all-order ε-expansion of one-loop two-point and of certain three-point functions was done explicitly by expanding the relevant hypergeometric functions. One-fold integral representations for general three- and four-point functions, as well as ways to get expansion terms of order ε for 3-point functions, were worked out in a recent paper [12]. Publications [9, 11, 12] also contain a comprehensive list of references on the subject.

However, in general, the required ε-expansion (including ε^2-terms) is not readily available for all the integrals needed in the hadronic heavy flavour production process. Also, the analytic continuation of the above mentioned hypergeometric functions in [9, 12] to the appropriate kinematical regions of validity is not always possible. This mainly concerns the four-point functions. In addition, it is more convenient to present results for the ε-expansion in terms of simpler special functions, in the form convenient for numerical evaluation. And finally, collecting together all the necessary scalar integrals needed for the derivation of tensor integrals entering the loop-by-loop contribution constitutes a first step in the difficult task of obtaining the NNLO corrections to heavy flavor hadroproduction cross section.

In our notation we shall remain very close to the notation introduced and used in [2,13]. In particular, we use dimensional regularization working in $n = 4 - 2\varepsilon$ dimensions, as e.g. in [13]. For the calculation of the NNLO virtual corrections to hadroproduction of heavy flavors one needs the same set of scalar master integrals as given in the Appendix A of [2] (the relevant set of master integrals is listed in Table 1). However, as explained above, knowledge of their singular and finite terms is not sufficient for the calculation of NNLO loop-by-loop corrections. For that purpose one needs to know the one-loop integrals up to $\mathcal{O}(\varepsilon^2)$ including also their imaginary parts which equally well contribute to the modulus squared of the one-loop amplitudes. The imaginary parts of the one-loop integrals are really needed only up to $\mathcal{O}(\varepsilon)$ since the highest singularity of the imaginary parts is only $\mathcal{O}(\varepsilon^{-1})$ compared to $\mathcal{O}(\varepsilon^{-2})$ for the real parts. We have nevertheless decided to include $\mathcal{O}(\varepsilon^2)$ results also for the imaginary parts which may be of interest in other applications. Consequently, in this paper we present the relevant expressions for all scalar integrals needed in the calculation of the NNLO loop-by-loop corrections to hadroproduction of heavy flavors. For reasons of comprehensiveness we have decided to include also the singular and finite (i.e. $\mathcal{O}(\varepsilon^0)$) parts of the scalar integrals in our presentation. They agree with the results of the real contributions presented in [2].

---

[1] In a more general setting the Laurent-series expansion of the scalar integrals is needed if the integration-by-parts technique [8] is employed. The reason is that the solution of the recursion relations induced by the integration-by-parts technique can bring in negative powers of ε.

A comment on the length of the formula expressions in our paper is appropriate. The untreated computer output of the integrations is generally quite lengthy. The hard work is to simplify these expressions. We have written semi–automatic computer codes that achieve the simplifications using known identities among polylogarithms and using a number of identities for the L–functions introduced in this paper which are derived in an Appendix.

The paper is organized as follows. In Sec. II we deal with one- and two-point functions. Sec. III contains our results on three-point functions, while in Sec. IV we present our results for the ε-expansion of the four-point functions. In Sec. V we give our summary and conclusions. We collect some technical material in three Appendixes. In Appendix A we discuss the Taylor series expansion around $p^2 = m^2$ for the self–energy insertion two–point function which is needed for the calculation of the heavy quark wave function renormalization constant. In Appendix B we define multiple polylogarithms and demonstrate that our analytical results can all be expressed in terms of multiple polylogarithms. In Appendix C we derive a number of identities for the so–called L–functions introduced in the main text. A judicious use of these identities has allowed us to considerably reduce the length of our final analytical results for the three– and four–point functions.

II. ONE- AND TWO-POINT FUNCTIONS

We start with the one-loop one-point function which is defined by

$$A(m) = \mu^{2\varepsilon} \int \frac{d^n q}{(2\pi)^n} \frac{1}{q^2 - m^2}. \tag{2.1}$$

Working in $n = 4 - 2\varepsilon$ dimensions, the expansion for this one-point scalar integral can be written in the general form:

$$A(m) = iC_\varepsilon(m^2) \frac{m^2}{\varepsilon(1-\varepsilon)} = iC_\varepsilon(m^2) m^2 \left\{ \frac{1}{\varepsilon} + 1 + \varepsilon + \varepsilon^2 + \mathcal{O}(\varepsilon^3) \right\}. \tag{2.2}$$

Table 1: List of one-, two-, three- and four-point massive one-loop functions calculated in this paper up to $\mathcal{O}(\varepsilon^2)$.

	Nomenclature of [2]	Our nomenclature	Novelty	Comments
1-point	$A(m)$	A	–	Re
2-point	$B(p_4 - p_2, 0, m)$	B_1	–	Re
	$B(p_3 + p_4, m, m)$	B_2	–	Re, Im
	$B(p_4, 0, m)$	B_3	–	Re
	$B(p_2, m, m)$	B_4	–	Re
	$B(p_3 + p_4, 0, 0)$	B_5	–	Re, Im
3-point	$C(p_4, p_3, 0, m, 0)$	C_1	new	Re, Im
	$C(p_4, -p_2, 0, m, m)$	C_2	new	Re
	$C(-p_2, p_4, 0, 0, m)$	C_3	–	Re
	$C(-p_2, -p_1, 0, 0, 0)$	C_4	–	Re, Im
	$C(-p_2, -p_1, m, m, m)$	C_5	–	Re, Im
	$C(p_3, p_4, m, 0, m)$	C_6	–	Re, Im
4-point	$D(p_4, -p_2, -p_1, 0, m, m, m)$	D_1	new	Re, Im
	$D(-p_2, p_4, p_3, 0, 0, m, 0)$	D_2	new	Re, Im
	$D(-p_2, p_4, -p_1, 0, 0, m, m)$	D_3	new	Re

where m is the internal loop mass and where we have defined

$$C_\varepsilon(m^2) \equiv \frac{\Gamma(1+\varepsilon)}{(4\pi)^2}\left(\frac{4\pi\mu^2}{m^2}\right)^\varepsilon . \tag{2.3}$$

The one-loop two-point functions are defined by [2]

$$B(q_1, m_1, m_2) = \mu^{2\varepsilon}\int \frac{d^n q}{(2\pi)^n}\frac{1}{(q^2 - m_1^2)[(q+q_1)^2 - m_2^2]}, \tag{2.4}$$

where the m_i $(i = 1, 2)$ can be either m or 0. In the denominators of the relevant functions we always imply the "causal" $+i\delta$ prescription to deal with singularities in pseudo-Euclidean space.

In what follows, we will always present our results for the scalar functions separately for the real and imaginary contributions. We introduce the Mandelstam-type variables

$$s \equiv (p_1+p_2)^2, \qquad t \equiv T - m^2 \equiv (p_1-p_3)^2 - m^2, \qquad u \equiv U - m^2 \equiv (p_2-p_3)^2 - m^2, \tag{2.5}$$

with the kinematical condition on external momenta being $p_1+p_2 = p_3+p_4$ (i.e. $s+t+u = 0$) and the on-shell conditions are $p_1^2 = p_2^2 = 0,\ p_3^2 = p_4^2 = m^2$. Note that the variables t and u defined in (2.5) are *not* the usual Mandelstam variables.

There are altogether five different two-point scalar functions B_i $(i = 1, 2, ..., 5)$ needed for hadronic heavy flavor production [2]. Again we choose to extract a common factor

$iC_\varepsilon(m^2)$, where $C_\varepsilon(m^2)$ is defined in (2.3). The coefficients of the ε-expansion are denoted by $B_i^{(j)}$, i.e. we write

$$B_i = i\, C_\varepsilon(m^2) \left\{ \frac{1}{\varepsilon} B_i^{(-1)} + B_i^{(0)} + \varepsilon B_i^{(1)} + \varepsilon^2 B_i^{(2)} + \mathcal{O}(\varepsilon^3) \right\}. \tag{2.6}$$

The ε-expansion of the two-point functions starts at ε^{-1}. It turns out that $B_i^{(-1)} = 1$ for all i. The first two-point function

$$B_1 \equiv B(p_4 - p_2, 0, m) \tag{2.7}$$

is real for our kinematics which can be seen by drawing the appropriate Feynman diagram for B_1 and applying the Landau-Cutkosky rules. The same statement holds true for the two-point functions B_3 and B_4 to be discussed later on. One has

$$\begin{aligned}
\mathrm{Re}\, B_1^{(-1)} &= 1, && (2.8)\\
\mathrm{Re}\, B_1^{(0)} &= 2 - \frac{t}{T} \ln \frac{-t}{m^2}, \\
\mathrm{Re}\, B_1^{(1)} &= 2\mathrm{Re}\, B_1^{(0)} + \frac{t}{T} \ln^2 \frac{-t}{m^2} + \frac{t}{T} \mathrm{Li}_2(\frac{T}{m^2}), \\
\mathrm{Re}\, B_1^{(2)} &= 2\mathrm{Re}\, B_1^{(1)} - \frac{t}{T} \left[\frac{1}{3} \ln^3 \frac{-t}{m^2} + 2\mathrm{Li}_3(\frac{T}{t}) + \mathrm{Li}_3(\frac{T}{m^2}) \right]; \\
\mathrm{Im}\, B_1^{(j)} &= 0. && (2.9)
\end{aligned}$$

The second scalar two-point function

$$B_2 \equiv B(p_3 + p_4, m, m) \tag{2.10}$$

has both real and imaginary parts. Defining $\beta = (1 - 4m^2/s)^{1/2}$ and $x = (1-\beta)/(1+\beta)$ we obtain

$$\begin{aligned}
\mathrm{Re}\, B_2^{(-1)} &= 1, && (2.11)\\
\mathrm{Re}\, B_2^{(0)} &= 2 + \beta \ln x, \\
\mathrm{Re}\, B_2^{(1)} &= \beta \left[\frac{4}{\beta} - 2\ln(\frac{s\beta^2}{m^2}) + \frac{1}{2}\ln^2(\frac{s\beta^2}{m^2}) + 4\ln(1-x) - 2\ln^2(1-x) - 4\zeta(2) - 2\mathrm{Li}_2(x) \right], \\
\mathrm{Re}\, B_2^{(2)} &= 2\mathrm{Re}\, B_2^{(1)} + \beta \left[4\zeta(2)\ln(\frac{s\beta^2}{m^2}) - \frac{1}{6}\ln^3(\frac{s\beta^2}{m^2}) + 4\zeta(2)\ln(1-x) - 2\ln x \ln^2(1-x) \right. \\
&\qquad \left. + \frac{4}{3}\ln^3(1-x) + 2\zeta(3) - 4\mathrm{Li}_3(1-x) - 2\mathrm{Li}_3(x) \right]; \\
\mathrm{Im}\, B_2^{(0)} &= \pi\beta, \qquad \mathrm{Im}\, B_2^{(1)} = \pi\beta \left[2 - \ln(\frac{s\beta^2}{m^2}) \right], && (2.12)\\
\mathrm{Im}\, B_2^{(2)} &= 2\mathrm{Im}\, B_2^{(1)} + \pi\beta \left[\frac{1}{2}\ln^2(\frac{s\beta^2}{m^2}) - 2\zeta(2) \right].
\end{aligned}$$

The remaining three two-point functions have a simple structure:

$$\begin{aligned}
B_3 \equiv B(p_4,0,m) &= iC_\varepsilon(m^2)\frac{1}{\varepsilon(1-2\varepsilon)} \\
&= iC_\varepsilon(m^2)\left\{\frac{1}{\varepsilon}+2+4\varepsilon+8\varepsilon^2+\mathcal{O}(\varepsilon^3)\right\}; \qquad (2.13)\\
B_4 \equiv B(p_2,m,m) &= iC_\varepsilon(m^2)\frac{1}{\varepsilon}; \qquad (2.14)\\
B_5 \equiv B(p_3+p_4,0,0) &= iC_\varepsilon(m^2)\frac{\Gamma^2(1-\varepsilon)}{\Gamma(2-2\varepsilon)}\left(-\frac{s+i\delta}{m^2}\right)^{-\varepsilon}\frac{1}{\varepsilon}. \qquad (2.15)
\end{aligned}$$

The results for B_3 and B_4 in (2.13), (2.14) are not separately listed in the standard format $B_i^{(j)}$ which can of course be read off from the relevant expressions (2.13), (2.14). The two-point function B_5 (2.15) has both real and imaginary parts:

$$\begin{aligned}
&\mathrm{Re}\, B_5^{(-1)} = 1, \qquad (2.16)\\
&\mathrm{Re}\, B_5^{(0)} = 2-\ln\frac{s}{m^2},\\
&\mathrm{Re}\, B_5^{(1)} = 2\mathrm{Re}\, B_5^{(0)} - 4\zeta(2) + \frac{1}{2}\ln^2\frac{s}{m^2},\\
&\mathrm{Re}\, B_5^{(2)} = 2\mathrm{Re}\, B_5^{(1)} + 4\zeta(2)\ln\frac{s}{m^2} - \frac{1}{6}\ln^3\frac{s}{m^2} - 2\zeta(3);
\end{aligned}$$

$$\begin{aligned}
&\mathrm{Im}\, B_5^{(0)} = \pi, \qquad \mathrm{Im}\, B_5^{(1)} = \pi\left[2-\ln\frac{s}{m^2}\right], \qquad (2.17)\\
&\mathrm{Im}\, B_5^{(2)} = 2\mathrm{Im}\, B_5^{(1)} + \pi\left[\frac{1}{2}\ln^2\frac{s}{m^2} - 2\zeta(2)\right].
\end{aligned}$$

We have done various checks on the above results. First of all, they were double-checked, i.e. the results were obtained by two independent calculations. Secondly, they were checked numerically by verifying that the original integrals (after Feynman parametrization and integrating out the loop momentum, and for B_1 and B_2 also expanding in ε) are equal numerically to the final integrals. We have also verified our results by extracting the relevant expressions from general formulae given in [9,10]. In particular, our coefficients (2.8) may be obtained from Eq. (10) of the first reference of [9] and then using Eq. (2.14) of [10]. Our result (2.11), (2.12) can be obtained from Eqs. (2.10) and (2.14) of [10]. Finally, our expressions (2.13) – (2.15) can also be obtained from Eqs. (10), (17) and (8), respectively, of the first reference of [9].

There is one more special case of the two-point integral which is needed for the calculation of a self-energy insertion into external massive fermion lines. This integral is used for the definition of the fermion mass and wave function renormalization constants in the on-shell renormalization scheme. This specific two-point function is given in the Appendix A of this paper.

III. THREE-POINT FUNCTIONS

The one-loop three-point functions are defined by [2]

$$C(q_1,q_2,m_1,m_2,m_3) = \mu^{2\varepsilon}\int \frac{d^nq}{(2\pi)^n}\frac{1}{(q^2-m_1^2)[(q+q_1)^2-m_2^2][(q+q_1+q_2)^2-m_3^2]}. \quad (3.18)$$

The three masses m_1, m_2 and m_3 come in various combinations of zero and nonzero masses where all nonzero masses are equal to m as before. There are six different types of three-point functions C_i ($i = 1, 2, ..., 6$) needed for our purposes [2]. They have both real and imaginary parts except for C_2 and C_3 which are real. Again, this can be seen from the Feynman diagrams representing C_2 and C_3 and applying the Landau-Cutkosky rules. Their ε-expansion is again written in the following universal format:

$$C_i = i\,C_\varepsilon(m^2)\left\{\frac{1}{\varepsilon^2}C_i^{(-2)}+\frac{1}{\varepsilon}C_i^{(-1)}+C_i^{(0)}+\varepsilon C_i^{(1)}+\varepsilon^2C_i^{(2)}+\mathcal{O}(\varepsilon^3)\right\}, \quad (3.19)$$

where the ε-expansion now starts at $\varepsilon^{(-2)}$. Note that the $C_i^{(-2)}$ are purely real.

It turns out that the $\mathcal{O}(\varepsilon^2)$ results for the three-point functions can no longer be presented in terms of classical polylogarithms but require a new class of functions given by the one-fold integral representations defined below. To write down our results in a short and convenient form, we introduce the following functions:

$$L_{\sigma_1\sigma_2\sigma_3}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) = \int_0^1 dy\frac{\ln(\alpha_1+\sigma_1y)\ln(\alpha_2+\sigma_2y)\ln(\alpha_3+\sigma_3y)}{\alpha_4+y}, \quad (3.20)$$

and

$$L_{\sigma_1}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) = \int_0^1 dy\frac{\ln(\alpha_1+\sigma_1y)\mathrm{Li}_2(\alpha_2+\alpha_3y)}{\alpha_4+y}. \quad (3.21)$$

Here the σ_i ($i = 1, 2, 3$) take values ±1 and the α_j's are either integers $\{1, 0, -1\}$ or else kinematical variables. The above L-functions arise naturally in our calculational framework [2]. They can all be expressed in terms of so-called multiple polylogarithms of maximum weight four [14] (see Appendix B for details). However, we choose to write our results in terms of the above single- and triple-index L-functions for several reasons. The results look simpler, e.g. they can be expressed as one-fold integrals of products of logarithms and dilogarithms, and are shorter. We have also found that the L-functions are much easier to evaluate numerically than the corresponding multiple polylogarithms (see [15] for relevant details).

There exist simple algebraic relations between these L-functions based on either symmetry relations regarding permutations of indices and change of integration variables or on relations based on integration-by-parts techniques. We describe them in Appendix C.

---

[2] As A. Davydychev informs us the functions analogous to our triple-index functions $L_{\sigma_1\sigma_2\sigma_3}$ also arise in the approach of [10] when one analytically continues their Eq. (3.2) for the order ε^2 terms.

In particular this means that our results on the three- and four-point functions can all be written in terms of the L_{-++} and L_{+++} variants of the triple–index $L_{\sigma_1\sigma_2\sigma_3}$ functions in Eq. (3.20), and of the L_+ variant of the single–index L_{σ_1} function of Eq. (3.21).

We start with the three-point function C_1 defined by

$$C_1 \equiv C(p_4, p_3, 0, m, 0).$$

One obtains

$$\begin{aligned}
\mathrm{Re}\, C_1^{(-2)} &= \mathrm{Re}\, C_1^{(-1)} = 0, \qquad\qquad (3.22)\\
\mathrm{Re}\, C_1^{(0)} &= \frac{1}{2s\beta}[\ln^2 x + 4\mathrm{Li}_2(-x) + 2\zeta(2)],\\
\mathrm{Re}\, C_1^{(1)} &= -\frac{1}{s\beta}\left[\frac{1}{6}\ln^3\frac{s}{m^2} + 2\ln\frac{s}{m^2}\ln(1-x)\ln x + \ln(1-x)\ln^2 x - 4\zeta(2)\ln\frac{s}{m^2}\right.\\
&+ 2\ln\frac{s}{m^2}\mathrm{Li}_2(x) + 5\zeta(3) - 4\mathrm{Li}_3(1-x) + 2\mathrm{Li}_3(x) + 2\mathrm{Li}_3(1-x^2)\\
&- \left. 8\mathrm{Li}_3(\frac{1}{1+x})\right],\\
\mathrm{Re}\, C_1^{(2)} &= \frac{1}{s\beta}\left[\frac{7}{48}\ln^4\frac{s}{m^2} - \frac{11}{24}\ln^3\frac{s}{m^2}\ \ln x - \frac{1}{4}\ln^2\frac{s}{m^2}\left(\ln^2(1-x) + 6\ln(1-x)\ln x\right.\right.\\
&+ \left. 5\ln^2 x\right) + \ln\frac{s}{m^2}\left(\frac{1}{3}\ln^3(1-x) - \frac{7}{2}\ln^2(1-x)\ln x + \frac{11}{24}\ln^3 x\right)\\
&- \frac{1}{3}\ln^3(1-x)\ln x - \frac{1}{4}\ln^2(1-x)\ln^2 x + \frac{5}{2}\ln(1-x)\ln^3 x + \frac{55}{48}\ln^4 x\\
&- \frac{7}{8}\left(3\ln^2\frac{s}{m^2} + 10\ln\frac{s}{m^2}\ln x + 16\ln(1-x)\ln x + 3\ln^2 x\right)\zeta(2)\\
&+ \frac{1}{2}\left(\ln\frac{s}{m^2} - 11\ln x\right)\zeta(3) + \frac{5}{2}\zeta(4) - 2\mathrm{Li}_2^2(-x) + \mathrm{Li}_2(x)\left(-\ln^2\frac{s}{m^2}\right.\\
&+ \left. 2\ln(1-x)\ln x + 7\ln^2 x - 2\ln\frac{s}{m^2}\left(\ln(1-x) - \ln x\right) - 10\,\zeta(2)\right)\\
&+ \mathrm{Li}_2(-x)\left(-\frac{13}{4}\ln^2\frac{s}{m^2} - 2\ln\frac{s}{m^2}\left(\ln(1-x) - \ln x\right) + 2\ln(1-x)\ln x\right.\\
&+ \left. \frac{31}{4}\ln^2 x - 6\,\zeta(2)\right) + 4\mathrm{Li}_3(1-x)\left(\ln\frac{s}{m^2} + 2\ln(1-x) - \ln x\right)\\
&- \mathrm{Li}_3(-x)\left(\frac{21}{2}\ln\frac{s}{m^2} + 20\ln(1-x) + \frac{29}{2}\ln x\right) - 4\mathrm{Li}_3(x)\left(\ln\frac{s}{m^2} + 2\ln(1-x)\right.\\
&+ \left. \frac{5}{2}\ln(x)\right) - \mathrm{Li}_3\left(\frac{1}{1+x}\right)\left(3\ln\frac{s}{m^2} + 4\ln(1-x) - 2\ln x\right) - 2\mathrm{Li}_3\left(1-x^2\right)\times\\
&\left(\ln\frac{s}{m^2} + 2\ln(1-x) - \ln x\right) - 4\,\mathrm{Li}_4(1-x) - 4\,\mathrm{Li}_4\left(\frac{-x}{1-x}\right) + 12\,\mathrm{Li}_4(-x)\\
&+ 28\mathrm{Li}_4(x) + 17\mathrm{Li}_4\left(\frac{1}{1+x}\right) - 17\mathrm{Li}_4\left(\frac{x}{1+x}\right) + 4\mathrm{Li}_4\left(1-x^2\right) + 4\mathrm{Li}_4\left(\frac{-x^2}{1-x^2}\right)
\end{aligned}$$

$$
\begin{aligned}
&- && L_{-++}\Big(1,x^{-1},x^{-1},0\Big)+L_{-++}\Big(1,x^{-1},x^{-1},x\Big)+2\,L_{-++}\Big(1,x,x^{-1},0\Big)\\
&- && 2\,L_{-++}\Big(1,x,x^{-1},x^{-1}\Big)-2\,L_{-++}\Big(1,x,x^{-1},x\Big)+L_{-++}(1,x,x,0)\\
&- && L_{-++}\Big(1,x,x,x^{-1}\Big)+2\,L_{-++}\Big(1+x^{-1},0,0,-1\Big)-2\,L_{-++}(1+x,0,0,-1)\\
&- && 2\,L_{-++}\Big(1+x^{-1},0,0,-1-x\Big)+2\,L_{-++}\Big(1+x,0,0,-1-x^{-1}\Big)\\
&+ && 2\,L_{+}\Big(0,0,-x^{-1},x\Big)-2\,L_{+}\Big(0,0,-x,x^{-1}\Big)-4\,L_{+}(0,0,-x,x)\\
&- && 4\,L_{+}\Big(0,-x^{-1},x^{-1},-1\Big)+4\,L_{+}\Big(0,-x^{-1},x^{-1},-1-x\Big)+4\,L_{+}(0,-x,x,-1)\\
&+ && 4\,L_{+}\Big(0,-x^{-1},x^{-1},-1-x^{-1}\Big)-4\,L_{+}(0,-x,x,-1-x)\\
&- && 4\,L_{+}\Big(0,-x,x,-1-x^{-1}\Big)-4\,L_{+}\Big(0,\frac{-x}{1-x},\frac{x}{1-x^2},-1\Big)\\
&+ && 4\,L_{+}\Big(0,\frac{-x}{1-x},\frac{x}{1-x^2},-1-x\Big)+4\,L_{+}\Big(0,\frac{-x}{1-x},\frac{x}{1-x^2},-1-x^{-1}\Big)\\
&+ && 2\,L_{+}\Big(x^{-1},0,-x^{-1},x\Big)+2\,L_{+}\Big(x^{-1},0,-x,x\Big)-3\,L_{+++}\Big(0,0,x^{-1},x\Big)\\
&- && L_{+++}\Big(0,0,x,x^{-1}\Big)+\frac{3}{2}\,L_{+++}\Big(0,x^{-1},x^{-1},x\Big)+3\,L_{+++}(0,x^{-1},x,x)\\
&+ && L_{+++}\Big(0,x,x^{-1},x^{-1}\Big)+\frac{1}{2}\,L_{+++}\Big(0,x,x,x^{-1}\Big)-2\,L_{+++}\Big(x^{-1},x,x^{-1},x\Big)\Big]\,;
\end{aligned}
$$

$$
\begin{aligned}
\mathrm{Im}\,C_1^{(-1)} &= 0, \qquad\qquad \mathrm{Im}\,C_1^{(0)}=\frac{\pi}{s\beta}\ln x, \qquad\qquad (3.23)\\
\mathrm{Im}\,C_1^{(1)} &= -\frac{\pi}{2s\beta}\left[2\ln\frac{s}{m^2}\ln x-4\ln(1-x)\ln x+\ln^2 x+4\zeta(2)-4\mathrm{Li}_2(x)\right],\\
\mathrm{Im}\,C_1^{(2)} &= \frac{\pi}{6s\beta}\left[3\ln^2\frac{s}{m^2}\ln x-12\ln\frac{s}{m^2}\ln(1-x)\ln x+3\ln\frac{s}{m^2}\ln^2 x-6\ln(1-x)\ln^2 x\right.\\
&+ \left.\ln^3 x-12\ln\frac{s}{m^2}\mathrm{Li}_2(x)-24\mathrm{Li}_3(1-x)-12\mathrm{Li}_3(x)+12\zeta(2)\ln\frac{s}{m^2}+12\zeta(3)\right].
\end{aligned}
$$

We have not been able to derive the corresponding result from the known general hypergeometric function that represents the above integral in [9]. On the other hand, for a general three-point function, an expression for the order ε-terms was obtained in [16] in terms of simple polylogarithms up to Li_3. However, we believe that the result Eq. (5.21) in [16] is not applicable to our case as one faces singularities resulting from vanishing denominators in the arguments of the relevant logarithms and polylogarithms. We have checked our final result numerically against the original two-fold and one-fold Feynman parameter integrals (after ε-expanding the corresponding integrand). This was done term by term for coefficients at the corresponding orders in ε. Although the result for the ε^2-coefficient looks lengthy, our final analytic results (3.22), (3.23) for the three-point function C_1 integrate out numerically very fast (in fraction of a second on a desktop computer for a chosen numerical point) and without any problems. In comparison, the numerical integration of the one-fold integral by Mathematica took eight times longer, and that of the two-fold integral even 200 times longer to evaluate. In addition, because of various branch cuts, the one- and two-fold integrals would only allow integrations in the complex plane of kinematical variables, while for the physical region they have severe problems.

The integral C_2 is real and finite:

$$C_2 \equiv C(p_4, -p_2, 0, m, m);$$

$$\begin{aligned}
\mathrm{Re}\, C_2^{(-2)} &= \mathrm{Re}\, C_2^{(-1)} = 0, && (3.24)\\
\mathrm{Re}\, C_2^{(0)} &= \frac{1}{t}\left[\zeta(2) - \mathrm{Li}_2(\frac{T}{m^2})\right],\\
\mathrm{Re}\, C_2^{(1)} &= \frac{1}{3t}\left[\ln^3\frac{-t}{m^2} + 6\ln\frac{-t}{m^2}\mathrm{Li}_2(\frac{T}{m^2}) + 9\zeta(3) - 6\mathrm{Li}_3(\frac{T}{t}) - 9\mathrm{Li}_3(\frac{T}{m^2})\right],\\
\mathrm{Re}\, C_2^{(2)} &= \frac{1}{24t}\left[-5\ln^4\frac{-t}{m^2} + 12\zeta(2)\ln^2\frac{-t}{m^2} + 12\zeta(2)\ln^2\frac{-T}{m^2} - 12\ln^2\frac{-t}{m^2}\mathrm{Li}_2(\frac{T}{m^2})\right.\\
&\quad -24\ln\frac{-t}{m^2}\ln\frac{-T}{m^2}\mathrm{Li}_2(\frac{T}{m^2}) - 24\zeta(3)\ln\frac{-t}{m^2} + 24\zeta(3)\ln\frac{-T}{m^2}\\
&\quad +24\ln\frac{-t}{m^2}\mathrm{Li}_3(\frac{T}{t}) + 24\ln\frac{-T}{m^2}\mathrm{Li}_3(\frac{T}{t}) + 24\ln\frac{-t}{m^2}\mathrm{Li}_3(\frac{T}{m^2})\\
&\quad +48\ln\frac{-T}{m^2}\mathrm{Li}_3(\frac{T}{m^2}) + 192\zeta(4) - 24\mathrm{Li}_4(\frac{m^2}{-t})\\
&\quad \left. +12L_{-++}(1, -\frac{m^2}{T}, -\frac{m^2}{T}, 0) - 12L_{-++}(\frac{t}{T}, 0, 0, -1)\right];\\
\mathrm{Im}\, C_2^{(j)} &= 0. && (3.25)
\end{aligned}$$

This result was checked numerically against the original double parametric representation (obtained after doing Feynman parametrization) of this integral expanded in powers of ε. We could not obtain similar expressions from known general results for this integral, as the ε-expansion of the relevant hypergeometric function is problematic. In addition, it turns out that the general result for the order ε-terms for the massive three-point function of [16] does not allow for a straightforward extraction of the corresponding expression for this particular case. More exactly, the equation $Q_3(y) = 0$ originating from the table in [16] (on page 608), does not have solutions for the relevant kinematics. In this sense, our expressions for the coefficients of the ε- and ε^2-terms for C_2 represent a new result.

The integration of the function C_3 defined by

$$C_3 \equiv C(-p_2, p_4, 0, 0, m)$$

requires the construction of a subtraction term since an ε-expansion of the relevant integrand does not straightforwardly lead to the desired ε-expansion of the integral. This is best illustrated in a simple example which nevertheless captures the essential idea of the subtraction method. Consider the integral

$$\int_0^1 dx\, x^{-1+\varepsilon} f(x, \varepsilon) \qquad (3.26)$$

where $f(x,\varepsilon)$ is an integrable function in the interval $[0,1]$ and has derivatives in ε. For the sake of the argument take $f(x,\varepsilon)$ to have a Laurent series expansion starting at the zeroth order in ε, i.e. $f(x,\varepsilon) = f^{(0)}(x) + \varepsilon f^{(1)}(x) + \ldots$. It is clear that expanding the integrand

$$x^{-1+\varepsilon} f(x,\varepsilon) = \frac{1}{x} f^{(0)}(x) + \varepsilon \left(\frac{\ln x}{x} f^{(0)}(x) + \frac{1}{x} f^{(1)}(x) \right) + \ldots$$

does not in general render the integral integrable. However, if we write

$$\int_0^1 dx x^{-1+\varepsilon} f(x,\varepsilon) = \int_0^1 dx \left(x^{-1+\varepsilon} \right)_+ f(x,\varepsilon) + f(0,\varepsilon) \int_0^1 dx x^{-1+\varepsilon} \tag{3.27}$$

the terms on the r.h.s. of (3.27) are now integrable. In (3.27) we have introduced a "plus" prescription

$$\int_0^1 dx \left(x^{-1+\varepsilon} \right)_+ f(x,\varepsilon) = \int_0^1 dx x^{-1+\varepsilon} (f(x,\varepsilon) - f(0,\varepsilon))$$

not unlike the "plus" prescription usually introduced when discussing parton splitting functions. The ε-expansion of the integral (3.26) can now be obtained since the first integrand on the r.h.s. of (3.27) can be expanded in ε and then be integrated term by term whereas the second integral can be computed in closed form. The task is then to find the appropriate subtraction terms for the integrals encountered in our calculation. This is required for the three-point function C_3 and the three four-point functions to be discussed in the next section.

As exemplified above we derive the subtraction terms by substituting the value of the integration variable (usually the lower or upper limit of integration) at which the given integrand diverges into the nonsingular part of the singular integrand. Adding and subtracting the subtraction term does all the job: e.g. the subtraction term contains all the poles in a given Feynman parameter but can be easily integrated due to its simpler analytic structure, while the rest of the integrand is now finite with respect to the same parameter and can therefore be integrated as well. When dealing with such a finite but complicated integration we often make use of the integration-by-parts method to evaluate and simplify our expressions.

Applying the subtraction method to the evaluation of the three-point function C_3 one obtains:

$$\begin{aligned}
\mathrm{Re}\, C_3^{(-2)} &= \frac{1}{2t}, \quad \mathrm{Re}\, C_3^{(-1)} = -\frac{1}{t} \ln \frac{-t}{m^2}, \quad \mathrm{Re}\, C_3^{(0)} = \frac{1}{t} \left[\ln^2 \frac{-t}{m^2} + \mathrm{Li}_2(\frac{T}{m^2}) \right], \\
\mathrm{Re}\, C_3^{(1)} &= -\frac{1}{t} \left[\frac{1}{3} \ln^3 \frac{-t}{m^2} + 2\mathrm{Li}_3(\frac{T}{t}) + \mathrm{Li}_3(\frac{T}{m^2}) \right], \qquad (3.28) \\
\mathrm{Re}\, C_3^{(2)} &= \frac{1}{3t} \left[\ln^4 \frac{-t}{m^2} - \ln^3 \frac{-t}{m^2} \ln \frac{-T}{m^2} - 3\zeta(2) \ln^2 \frac{-t}{m^2} + 6\zeta(3) \ln \frac{-t}{m^2} - 6\zeta(4) \right. \\
&\left. \quad -3\mathrm{Li}_4(\frac{T}{m^2}) + 6\mathrm{Li}_4(\frac{m^2}{-t}) - 6\mathrm{Li}_4(\frac{T}{t}) \right]; \\
\mathrm{Im}\, C_3^{(j)} &= 0. \qquad (3.29)
\end{aligned}$$

Note that one can obtain corresponding expressions in terms of generalized Nielsen polylogarithms from Eq. (27) of [9]. The corresponding hypergeometric function of three variables Φ_1 can be reduced to a hypergeometric function ${}_2F_1$ of one variable and one can then use Eq. (2.14) of [10] to get the relevant ε-expansion. We have verified agreement with [9] analytically up to $\mathcal{O}(\varepsilon)$. The agreement for the ε^2-terms was verified numerically.

The three-point function C_4 has a closed form solution:

$$C_4 \equiv C(-p_2, -p_1, 0, 0, 0) = \frac{iC_\varepsilon(m^2)}{s} \frac{\Gamma^2(-\varepsilon)}{\Gamma(1-2\varepsilon)} \left(-\frac{s+i\delta}{m^2} \right)^{-\varepsilon} \tag{3.30}$$

which is straightforward to obtain. For the ε-expansion of the real and imaginary parts of (3.30) we get:

$$\begin{aligned}
\operatorname{Re} C_4^{(-2)} &= \frac{1}{s}, \quad \operatorname{Re} C_4^{(-1)} = -\frac{1}{s} \ln \frac{s}{m^2}, \quad \operatorname{Re} C_4^{(0)} = \frac{1}{2s} \left[\ln^2 \frac{s}{m^2} - 8\zeta(2) \right], \\
\operatorname{Re} C_4^{(1)} &= \frac{1}{s} \left[4\zeta(2) \ln \frac{s}{m^2} - \frac{1}{6} \ln^3 \frac{s}{m^2} - 2\zeta(3) \right], \\
\operatorname{Re} C_4^{(2)} &= \frac{1}{s} \left[\frac{1}{24} \ln^4 \frac{s}{m^2} - 2\zeta(2) \ln^2 \frac{s}{m^2} + 2\zeta(3) \ln \frac{s}{m^2} + 9\zeta(4) \right];
\end{aligned} \tag{3.31}$$

$$\begin{aligned}
\operatorname{Im} C_4^{(-1)} &= \frac{\pi}{s}, \quad \operatorname{Im} C_4^{(0)} = -\frac{\pi}{s} \ln \frac{s}{m^2}, \\
\operatorname{Im} C_4^{(1)} &= \frac{\pi}{2s} \left[\ln^2 \frac{s}{m^2} - 4\zeta(2) \right], \\
\operatorname{Im} C_4^{(2)} &= \frac{\pi}{s} \left[2\zeta(2) \ln \frac{s}{m^2} - \frac{1}{6} \ln^3 \frac{s}{m^2} - 2\zeta(3) \right].
\end{aligned} \tag{3.32}$$

For the fifth three-point integral C_5 defined by

$$C_5 \equiv C(-p_2, -p_1, m, m, m)$$

we first obtain a one-fold integral representation similar to Eq. (3.13) of [10]. As before, the main difficulty is the derivation of the coefficient for the ε^2-term. The corresponding coefficient has a complicated singularity structure as well as two branch points on its integration path. Therefore, in order to analytically separate the real and imaginary parts for our final result, we have divided the integration regions for the relevant terms into three parts. After analytical integration these terms are free of numerical instabilities and converge very fast. One obtains

$$\begin{aligned}
\operatorname{Re} C_5^{(-2)} &= \operatorname{Re} C_5^{(-1)} = 0, \\
\operatorname{Re} C_5^{(0)} &= \frac{1}{2s} \left[\ln^2 x - 6\zeta(2) \right], \\
\operatorname{Re} C_5^{(1)} &= \frac{1}{2s} \left[\frac{1}{3} \ln^3 x - 8\zeta(2) \ln x + 4 \ln x \operatorname{Li}_2(x) - 6\zeta(3) - 8\operatorname{Li}_3(x) \right],
\end{aligned} \tag{3.33}$$

$$
\begin{aligned}
\operatorname{Re} C_5^{(2)} &= \frac{1}{2s}\Big[4\ln x\ln^3(1-x)-\frac{9}{2}\ln^2 x\ln^2(1-x)-\frac{1}{3}\ln^3 x\ln(1-x)+\frac{1}{12}\ln^4 x\\
&\quad -\zeta(2)\left(3\ln^2 x-10\ln x\ln(1-x)+6\ln^2(1-x)\right)-\ln^2 x\,\mathrm{Li}_2(-x)\\
&\quad +2\left(\ln^2 x-6\ln x\ln(1-x)+3\ln^2(1-x)-6\zeta(2)\right)\mathrm{Li}_2(x)+10\zeta(3)\ln x\\
&\quad -6\zeta(3)\ln(1-x)+2\left(\ln x+3\ln(1-x)\right)\mathrm{Li}_3(x)+2\ln x\,\mathrm{Li}_3(-x)\\
&\quad -12\left(\ln x-\ln(1-x)\right)\mathrm{Li}_3(1-x)+\frac{45}{2}\zeta(4)-L_{-++}\left(1,0,0,\frac{1}{-1+x}\right)\\
&\quad +L_{-++}\left(1,0,0,\frac{x}{1-x}\right)-L_{+++}\left(0,0,\frac{1-x}{x},-1\right)+L_{+++}\left(0,0,\frac{1-x}{x},\frac{1}{x}\right)\\
&\quad -L_{+++}\left(0,\frac{1-x}{x},\frac{1-x}{x},-1\right)+L_{+++}\left(0,\frac{1-x}{x},\frac{1-x}{x},\frac{1}{x}\right)\Big];\\
\operatorname{Im} C_5^{(-1)} &= 0,\qquad \operatorname{Im} C_5^{(0)}=\frac{\pi}{s}\ln x, \qquad\qquad (3.34)\\
\operatorname{Im} C_5^{(1)} &= \frac{\pi}{2s}\left[\ln^2 x-4\zeta(2)+4\mathrm{Li}_2(x)\right],\\
\operatorname{Im} C_5^{(2)} &= \frac{\pi}{6s}\big[-12\ln x\ln^2(1-x)+\ln^3 x-12(\ln x-2\ln(1-x))(\zeta(2)-\mathrm{Li}_2(x))\\
&\quad +12\zeta(3)-12\mathrm{Li}_3(x)-24\mathrm{Li}_3(1-x)\big].
\end{aligned}
$$

Explicit result for this integral was given very recently in Eq. (4.4) of [11]. We have checked agreement with [11] analytically up to $\mathcal{O}(\varepsilon)$. The agreement for the ε^2-terms was verified numerically.

Finally, we write down real and imaginary parts for the last required three-point function C_6 defined by

$$
C_6 \equiv C(p_3, p_4, m, 0, m).
$$

One has

$$
\begin{aligned}
\operatorname{Re} C_6^{(-2)} &= 0,\qquad \operatorname{Re} C_6^{(-1)}=\frac{1}{s\beta}\ln x, \qquad\qquad (3.35)\\
\operatorname{Re} C_6^{(0)} &= \frac{1}{2s\beta}\left[-4\ln x\ln(1-x)+\ln^2 x-8\zeta(2)-4\mathrm{Li}_2(x)\right],\\
\operatorname{Re} C_6^{(1)} &= \frac{1}{6s\beta}\big[-6\ln^2 x\ln(1-x)+\ln^3 x-24\zeta(2)\ln x+72\zeta(2)\ln(1-x)+12\zeta(3)\\
&\quad -12\mathrm{Li}_3(x)-24\mathrm{Li}_3(1-x)\big],\\
\operatorname{Re} C_6^{(2)} &= \frac{1}{24s\beta}\Big[-16\ln x\ln^3(1-x)+24\ln^2 x\ln^2(1-x)-8\ln^3 x\ln(1-x)+\ln^4 x\\
&\quad +4\ln^4(1-x)+192\zeta(2)\ln x\ln(1-x)-48\zeta(2)\ln^2 x-240\zeta(2)\ln^2(1-x)\\
&\quad -48\zeta(3)\ln x+120\zeta(4)+48\mathrm{Li}_4(x)+96\mathrm{Li}_4\Big(\frac{-x}{1-x}\Big)+96\mathrm{Li}_4(1-x)\Big];\\
\operatorname{Im} C_6^{(-1)} &= \frac{\pi}{s\beta},\qquad \operatorname{Im} C_6^{(0)}=\frac{\pi}{s\beta}\left[\ln x-2\ln(1-x)\right], \qquad\qquad (3.36)
\end{aligned}
$$

$$
\begin{aligned}
\operatorname{Im} C_6^{(1)} &= \frac{\pi}{6s\beta}\left[-12\ln x\ln(1-x)+3\ln^2 x+12\ln^2(1-x)-12\zeta(2)\right], \\
\operatorname{Im} C_6^{(2)} &= \frac{\pi}{6s\beta}\left[12\ln x\ln^2(1-x)-6\ln^2 x\ln(1-x)+\ln^3 x-8\ln^3(1-x)-12\zeta(2)\ln x\right. \\
&\quad \left.+24\zeta(2)\ln(1-x)-12\zeta(3)\right].
\end{aligned}
$$

Corresponding results for C_6 may be obtained from Eqs. (3.5), (3.7), (2.10) and (2.14) of [10]. We have done an order by order numerical comparisons for the coefficients of the ε- and ε^2-terms, while other terms can be easily compared analytically. We obtain exact agreement.

We mention that we have checked all our analytical results for the three–point functions against numerical results provided to us by M.M. Weber [17] (see also [18]). We have found agreement.

IV. FOUR-POINT FUNCTIONS

The scalar four-point one-loop integrals with one, two or three heavy quarks running in the loop are the most difficult to evaluate. The one-loop four-point functions are defined by [2]

$$
\begin{aligned}
&D(q_1,q_2,q_3,m_1,m_2,m_3,m_4) = \qquad (4.37)\\
&\quad \mu^{2\varepsilon}\int\frac{d^nq}{(2\pi)^n}\frac{1}{(q^2-m_1^2)[(q+q_1)^2-m_2^2][(q+q_1+q_2)^2-m_3^2][(q+q_1+q_2+q_3)^2-m_4^2]}.
\end{aligned}
$$

As before, the $+i\delta$ terms in the denominators have not been written out. Again, there is only one internal mass scale for our purposes.

For heavy flavor production one needs three different types of four-point functions D_i $(i=1,2,3)$ which are expanded as

$$
D_i = i\,C_\varepsilon(m^2)\left\{\frac{1}{\varepsilon^2}D_i^{(-2)}+\frac{1}{\varepsilon}D_i^{(-1)}+D_i^{(0)}+\varepsilon D_i^{(1)}+\varepsilon^2 D_i^{(2)}+\mathcal{O}(\varepsilon^3)\right\}. \qquad (4.38)
$$

Again the coefficient of the most singular part of the four-point functions is purely real, i.e. $\operatorname{Im} D_i^{(-2)}=0$.

Before we give our results for the four-point functions it is necessary to discuss some general technical features. After applying Feynman parametrization, we are left with a three-fold parametric integral for the four-point functions:

$$
D = iC_\varepsilon(m^2)(1+\varepsilon)(m^2)^{-2}\int_0^1 dx_1dx_2dx_3\; x_1^2x_2K^{-2-\varepsilon}, \qquad (4.39)
$$

with the kernel K given by

$$
\begin{aligned}
m^2K &= -abq_1^2-ac(q_1+q_2)^2-ad(q_1+q_2+q_3)^2-bcq_2^2 \\
&\quad -bd(q_2+q_3)^2-cdq_3^2+am_1^2+bm_2^2+cm_3^2+dm_4^2-i\delta, \qquad (4.40)
\end{aligned}
$$

where the m_i $(i = 1, \ldots, 4)$ can be either m or 0. The set of parameters $\{a, b, c, d\}$ above corresponds to an arbitrary choice from the set of original parameters $\{x_1x_2x_3,\ x_1x_2(1-x_3),\ x_1(1-x_2),\ 1-x_1\}$. For each particular four-point function we make a judicious choice of Feynman parameters which enables us to get the most convenient kernel for the subsequent integrations.



FIG. 2. Massive box D_1 with three massive propagators.

First we consider the four-point function D_1 with three massive propagators shown in Fig. 2 which is defined by

$$D_1 \equiv D(p_4, -p_2, -p_1, 0, m, m, m).$$

Substitution of the corresponding values of momenta and masses for D_1 into the expression for our kernel (4.40) gives

$$K = ac\tilde{t} - bd\tilde{s} + (1-a)^2 - i\delta, \tag{4.41}$$

where we have introduced positive valued dimensionless variables

$$\tilde{s} \equiv \frac{s}{m^2}, \qquad \tilde{t} \equiv -\frac{t}{m^2}. \tag{4.42}$$

The kinematical conditions $\tilde{s} \geq 4$, $\tilde{t} \geq 1$, $\tilde{s} \geq \tilde{t}$ constrain the allowable region of phase space for the present physical 2 → 2 process. Our choice for the parameters $\{a, b, c, d\}$ is $\{1-x_1,\ x_1x_2x_3,\ x_1x_2(1-x_3),\ x_1(1-x_2)\}$. For D_1, the integration of the corresponding integrand over x_3 results in two terms:

$$I^{D_1}_{x_1x_2} = -\frac{x_1^{-\varepsilon}\left[x_1 + \tilde{t}(1-x_1)x_2\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{s}x_1(1-x_2) + \tilde{t}(1-x_1)]}, \tag{4.43}$$

$$II^{D_1}_{x_1x_2} = \frac{x_1^{-1-2\varepsilon}\left[1 - \tilde{s}x_2(1-x_2) - i\delta\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{s}x_1(1-x_2) + \tilde{t}(1-x_1)]}, \tag{4.44}$$

which then have to be integrated over the remaining parameters x_1 and x_2. Eqs. (4.43) and (4.44) correspond to the indefinite integral (or primitive) evaluated at the upper and lower limit of x_3, respectively. The term $I^{D_1}_{x_1x_2}$ in (4.43) does not change sign on the integration path, e.g. does not have a branch cut in the interval $[0,1]$ for both variables x_1 and x_2. Consequently, it does not give an imaginary contribution and it is thus safe to drop the $i\delta$ shift in $I^{D_1}_{x_1x_2}$. Furthermore, since $I^{D_1}_{x_1x_2}$ does not have poles in ε, we expand it up to ε^2 and straightforwardly integrate over the second variable x_2 to obtain $I^{D_1}_{x_1}$. Concerning the second term $II^{D_1}_{x_1x_2}$ in (4.44), one can see that there is a branch cut for the variable x_2 in its numerator as well as a divergence due to the factor $x_1^{-1-2\varepsilon}$ at the lower limit of the integration $x_1 = 0$ (we have dropped the $i\delta$ shift in the denominator as it does not affect our further calculation). At this point we introduce a subtraction term for $II^{D_1}_{x_1x_2}$ in the simplest possible way: we set $x_1 = 0$ in $II^{D_1}_{x_1x_2}$ everywhere except for the divergent term $x_1^{-1-2\varepsilon}$. This results in the following subtraction term:

$$II^{D_1,s}_{x_1x_2} = \frac{x_1^{-1-2\varepsilon}\,[1-\tilde{s}x_2(1-x_2)-i\delta]^{-1-\varepsilon}}{(1+\varepsilon)\tilde{t}}, \tag{4.45}$$

which, in the framework of the dimensional regularization scheme, integrates over x_1 to

$$II^{D_1,s}_{x_2} = -\frac{[1-\tilde{s}x_2(1-x_2)-i\delta]^{-1-\varepsilon}}{2\varepsilon(1+\varepsilon)\tilde{t}}. \tag{4.46}$$

Then we expand the above expression up to ε^2 and reexpress the argument of subsequent logarithms as

$$1-\tilde{s}x_2(1-x_2)-i\delta = \frac{(x_2-x_2^{(0)})(x_2-1+x_2^{(0)})}{x_2^{(0)}(1-x_2^{(0)})}, \tag{4.47}$$

with

$$x_2^{(0)} = \frac{1+\sqrt{1-4/\tilde{s}}}{2} + i\delta = \frac{1+\beta}{2} + i\delta \tag{4.48}$$

A final integration of the subsequent series can be done analytically in the complex plane and its result is expressed in terms of logarithms and classical polylogarithms up to Li_4. Analytic continuation of the result for $\delta \to 0$ is then straightforward.

Lastly, we calculate the finite difference

$$\Delta II^{D_1}_{x_1x_2} = II^{D_1}_{x_1x_2} - II^{D_1,s}_{x_1x_2} = \frac{x_1^{-1-2\varepsilon}\left(\tilde{t}-\tilde{s}(1-x_2)\right)[1-\tilde{s}x_2(1-x_2)-i\delta]^{-1-\varepsilon}}{(1+\varepsilon)\,\tilde{t}\,[\tilde{s}x_1(1-x_2)+\tilde{t}(1-x_1)]}$$

by again expanding the difference up to ε^2 and using (4.47) for the arguments of the logarithms. Then we first integrate over the variable x_2, leading to a reduction of the integrand to simple fractions w.r.t. x_2. In this way we avoid spurious poles in the remaining integral which would otherwise arise in case of integration over x_1 first.

To complete the derivation of the first four-point (box) integral, we combine the two terms

$$I^{D_1}_{x_1} + \Delta II^{D_1}_{x_1}$$

and perform the last integration over the variable x_1.

At this point we would like to comment on some technical details of our calculation which are used throughout this work. For instance, the integrand for the last integration contains expressions such as

$$f(x_1) \cdot \mathrm{Li}_{2,3}\left(\frac{a_1 x_1^2 + a_2 x_1 + a_3}{a_4 x_1^2 + a_5 x_1 + a_6}\right), \tag{4.49}$$

where $f(x_1)$ is a rational function or a product of a rational function and a logarithm. Using recursively the method of integration-by-parts as much as necessary we render their arguments to be linear functions of x_1. In addition, in the case of Li_3, we can reduce the weight of Li_3 by one. At the same time, the sources of imaginary contributions are transferred into logarithms (or remain in Li_2's and Li_3's with arguments that are independent of the integration variable). Finally, performing the last integration and adding up all the relevant contributions we arrive at the result for the box integral with three massive lines, containing polylogs up to Li_4 and the single- and triple-index L-functions introduced in Eqs. (3.20) and (3.21).

In order to keep our results at reasonable length we introduce the abbreviations

$$\begin{aligned}
&z_3 \equiv (s + 2t + s\beta)/2, \qquad z_4 \equiv (s + 2t - s\beta)/2, \\
&z_5 \equiv (2m^2 + t + t\beta)/2, \qquad z_6 \equiv (2m^2 + t - t\beta)/2, \\
&l_s \equiv \ln\frac{s}{m^2}, \quad l_t \equiv \ln\frac{-t}{m^2}, \quad l_T \equiv \ln\frac{-T}{m^2}, \quad l_x \equiv \ln x, \\
&l_\beta \equiv \ln\beta, \quad l_{z3} \equiv \ln\frac{z_3}{m^2}, \quad l_{z4} \equiv \ln\frac{-z_4}{m^2}
\end{aligned} \tag{4.50}$$

and obtain:

$$\begin{aligned}
\mathrm{Re}\, D_1^{(-2)} &= 0, \qquad \mathrm{Re}\, D_1^{(-1)} = \ln x/(st\beta), \\
\mathrm{Re}\, D_1^{(0)} &= -\left[2\ln x \ln(-t\beta/m^2) + 2\mathrm{Li}_2(x) - 2\mathrm{Li}_2(-x) + 3\zeta(2)\right]/(st\beta), \\
\mathrm{Re}\, D_1^{(1)} &= \frac{1}{st\beta}\left[\frac{l_s^3}{12} - \frac{l_s^2\, l_x}{2} - \frac{5\, l_x^3}{6} - l_x^2\, l_{z3} + l_x^2\, l_{z4} - l_x\, l_{z4}^2 - \frac{l_t^3}{3} + l_t^2\,(3\, l_x + l_{z4}) - \right. \\
&\quad l_t\, \left(l_T^2 + l_x^2 - l_{z3}^2 + 2l_T\,(l_x + l_{z3} - l_{z4}) + l_{z4}^2 + l_x\,(2\, l_{z3} - 4\, l_\beta)\right) + l_x\, l_\beta^2 - l_\beta^3 + \\
&\quad l_s\, \left(l_t\, l_x + l_x^2/4 + l_x\, (l_{z3} + l_{z4}) - l_\beta^2\right) + (-l_s + 10l_t - l_x + 6l_\beta)\, \zeta(2) - \\
&\quad 5\zeta(3) + 4\mathrm{Li}_2(x)\, l_t - 2\mathrm{Li}_2\left(\frac{m^2\, x}{-T}\right) l_t + 2\mathrm{Li}_2\left(\frac{T}{z_3}\right) l_t - 2\mathrm{Li}_2\left(\frac{m^2}{z_5}\right) l_x - \\
&\quad 2\mathrm{Li}_2\left(\frac{-t\,(1-\beta)}{2\, m^2}\right) l_x - 2\mathrm{Li}_3(-x) - 2\mathrm{Li}_3\left(\frac{-x^2}{1-x^2}\right) - 2\mathrm{Li}_3\left(\frac{z_3}{t}\right) + \\
&\quad 2\mathrm{Li}_3\left(\frac{z_4}{t}\right) + 4\mathrm{Li}_3\left(\frac{1-\beta}{2}\right) + 2\mathrm{Li}_3\left(\frac{m^2\,(1-\beta)}{2\, z_5}\right) - 2\mathrm{Li}_3\left(\frac{-t\,(1-\beta)}{2\, z_5}\right) + \\
&\quad \left. 8\mathrm{Li}_3\left(\frac{-1+\beta}{2\,\beta}\right) - 2\mathrm{Li}_3\left(\frac{2\, z_6}{m^2(1+\beta}\right) + 2\mathrm{Li}_3\left(\frac{2\, z_6}{-t(1+\beta)}\right)\right],
\end{aligned} \tag{4.51}$$

$$
\begin{aligned}
\mathrm{Re}\, D_1^{(2)} \quad = \quad & \frac{1}{st\beta}\Bigg[-\frac{5\, l_s^4}{64} - \frac{43}{24}\, l_s^3\, l_t - \frac{7}{4}\, l_s^2\, l_t^2 + \frac{2}{3}\, l_s\, l_t^3 + \frac{5\, l_t^4}{12} - \frac{11}{24}\, l_s^3\, l_T - \frac{3}{2}\, l_s^2\, l_t\, l_T - \\
& \frac{5}{2}\, l_s\, l_t^2\, l_T - \frac{4}{3}\, l_t^3\, l_T - \frac{5}{16}\, l_s^2\, l_T^2 + \frac{1}{4}\, l_s\, l_t\, l_T^2 - \frac{1}{2}\, l_s\, l_T^3 + l_t\, l_T^3 + \frac{5}{8}\, l_s^3\, l_x - \frac{3}{8}\, l_s^2\, l_t\, l_x - \\
& 4\, l_s\, l_t^2\, l_x - l_t^3\, l_x - \frac{7}{8}\, l_s^2\, l_T\, l_x + \frac{3}{2}\, l_s\, l_t\, l_T\, l_x - \frac{3}{2}\, l_t^2\, l_T\, l_x + \frac{3}{8}\, l_s\, l_T^2\, l_x + \frac{13}{4}\, l_t\, l_T^2\, l_x - \\
& \frac{1}{2} l_T^3\, l_x - \frac{13}{32}\, l_s^2\, l_x^2 + \frac{1}{8}\, l_s\, l_t\, l_x^2 - \frac{1}{4}\, l_t^2\, l_x^2 + \frac{17}{8}\, l_s\, l_T\, l_x^2 + \frac{1}{2}\, l_t\, l_T\, l_x^2 + \frac{7}{16}\, l_T^2\, l_x^2 + \frac{1}{2} l_s\, l_x^3 + \\
& \frac{3}{8}\, l_t\, l_x^3 + \frac{17}{24}\, l_T\, l_x^3 + \frac{7\, l_x^4}{64} + \frac{29}{48}\, l_s^3\, l_{z3} + 3 l_s^2\, l_t\, l_{z3} + \frac{5}{2}\, l_s\, l_t^2\, l_{z3} - \frac{3}{2}\, l_t^3\, l_{z3} - \frac{3}{4}\, l_s^2\, l_T\, l_{z3} + \\
& \frac{17}{2}\, l_s\, l_t\, l_T\, l_{z3} + \frac{5}{2}\, l_t^2\, l_T\, l_{z3} + 2\, l_t\, l_T^2\, l_{z3} - \frac{19}{16}\, l_s^2\, l_x\, l_{z3} + l_s\, l_t\, l_x\, l_{z3} + \frac{5}{2}\, l_t^2\, l_x\, l_{z3} + \\
& \frac{9}{2}\, l_s\, l_T\, l_x\, l_{z3} - \frac{1}{2}\, l_t\, l_T\, l_x\, l_{z3} - \frac{1}{2}\, l_T^2\, l_x\, l_{z3} + \frac{13}{16}\, l_s\, l_x^2\, l_{z3} + l_t\, l_x^2\, l_{z3} - \frac{3}{4}\, l_T\, l_x^2\, l_{z3} - \\
& \frac{11}{48}\, l_x^3\, l_{z3} - \frac{1}{8}\, l_s^2\, l_{z3}^2 - 4\, l_s\, l_t\, l_{z3}^2 + \frac{1}{2}\, l_t^2\, l_{z3}^2 + l_s\, l_T\, {l_{z3}}^2 - 5\, l_t\, l_T\, l_{z3}^2 - \frac{3}{4}\, l_s\, l_x\, l_{z3}^2 + \\
& l_t\, l_x\, l_{z3}^2 - 2\, l_T\, l_x\, l_{z3}^2 - \frac{1}{8}\, l_x^2\, l_{z3}^2 + l_t\, l_{z3}^3 + \frac{61}{48}\, l_s^3\, l_{z4} + \frac{27}{4}\, l_s^2\, l_t\, l_{z4} + \frac{1}{2}\, l_s\, l_t^2\, l_{z4} - \\
& \frac{13}{6}\, l_t^3\, l_{z4} + \frac{9}{4}\, l_s^2\, l_T\, l_{z4} - \frac{1}{2}\, l_s\, l_t\, l_T\, l_{z4} + \frac{5}{2}\, l_t^2\, l_T\, l_{z4} + l_s\, l_T^2\, l_{z4} - \frac{5}{2}\, l_t\, l_T^2\, l_{z4} + \\
& \frac{9}{16}\, l_s^2\, l_x\, l_{z4} + \frac{11}{2}\, l_s\, l_t\, l_x\, l_{z4} + \frac{3}{2}\, l_t^2\, l_x\, l_{z4} - \frac{3}{2}\, l_s\, l_T\, l_x\, l_{z4} - \frac{5}{2}\, l_t\, l_T\, l_x\, l_{z4} + \frac{1}{2} l_T^2\, l_x\, l_{z4} - \\
& \frac{11}{16}\, l_s\, l_x^2\, l_{z4} - \frac{1}{4}\, l_t\, l_x^2\, l_{z4} - \frac{9}{4}\, l_T\, l_x^2\, l_{z4} - \frac{5}{16}\, l_x^3\, l_{z4} - 2\, l_s^2\, l_{z3}\, l_{z4} - 6\, l_s\, l_t\, l_{z3}\, l_{z4} - \\
& l_t^2\, l_{z3}\, l_{z4} - 3 l_s\, l_T\, l_{z3}\, l_{z4} - 6 l_t\, l_T\, l_{z3}\, l_{z4} - 4 l_t\, l_x\, l_{z3}\, l_{z4} - 3 l_T\, l_x\, l_{z3}\, l_{z4} - \frac{1}{2} l_x^2\, l_{z3}\, l_{z4} + \\
& l_s\, l_{z3}^2\, l_{z4} + \frac{5}{2}\, l_t\, l_{z3}^2\, l_{z4} + l_x\, l_{z3}^2\, l_{z4} - \frac{77}{16}\, l_s^2\, l_{z4}^2 - \frac{13}{2}\, l_s\, l_t\, l_{z4}^2 + 3\, l_t^2\, l_{z4}^2 - \frac{5}{4}\, l_s\, l_T\, l_{z4}^2 + \\
& 2\, l_t\, l_T\, l_{z4}^2 + \frac{1}{4}\, l_T^2\, l_{z4}^2 - \frac{33}{8}\, l_s\, l_x\, l_{z4}^2 - \frac{5}{2}\, l_t\, l_x\, l_{z4}^2 + \frac{7}{4}\, l_T\, l_x\, l_{z4}^2 + \frac{15}{16}\, l_x^2\, l_{z4}^2 + 3\, l_s\, l_{z3}\, l_{z4}^2 + \\
& \frac{7}{2}\, l_t\, l_{z3}\, l_{z4}^2 + 3\, l_T\, l_{z3}\, l_{z4}^2 + 2\, l_x\, l_{z3}\, l_{z4}^2 - l_{z3}^2\, l_{z4}^2 + \frac{71}{12}\, l_s\, l_{z4}^3 + \frac{2}{3}\, l_t\, l_{z4}^3 - \frac{5}{6}\, l_T\, l_{z4}^3 + \\
& \frac{31}{12}\, l_x\, l_{z4}^3 - 2 l_{z3}\, l_{z4}^3 - \frac{49}{24} l_{z4}^4 - \frac{13}{24}\, l_s^3\, l_\beta - \frac{7}{2}\, l_s^2\, l_t\, l_\beta - 2\, l_s\, l_t^2\, l_\beta - \frac{5}{3}\, l_t^3\, l_\beta - \frac{9}{4}\, l_s^2\, l_T\, l_\beta - \\
& 3\, l_s\, l_t\, l_T\, l_\beta - 4\, l_t^2\, l_T\, l_\beta + \frac{13}{8}\, l_s^2\, l_x\, l_\beta - 2\, l_s\, l_t\, l_x\, l_\beta - 5\, l_t^2\, l_x\, l_\beta - \frac{1}{2}\, l_s\, l_T\, l_x\, l_\beta - \\
& 3\, l_t\, l_T\, l_x\, l_\beta + l_T^2\, l_x\, l_\beta - \frac{9}{8}\, l_s\, l_x^2\, l_\beta + \frac{1}{2}\, l_t\, l_x^2\, l_\beta + \frac{7}{4}\, l_T\, l_x^2\, l_\beta + \frac{17}{24}\, l_x^3\, l_\beta + \frac{9}{4}\, l_s^2\, l_{z3}\, l_\beta + \\
& 4\, l_s\, l_t\, l_{z3}\, l_\beta + 4\, l_t^2\, l_{z3}\, l_\beta + 2\, l_s\, l_T\, l_{z3}\, l_\beta + 6\, l_t\, l_T\, l_{z3}\, l_\beta - \frac{5}{2}\, l_s\, l_x\, l_{z3}\, l_\beta + 4\, l_t\, l_x\, l_{z3}\, l_\beta + \\
& 4 l_T\, l_x\, l_{z3}\, l_\beta + \frac{5}{4} l_x^2\, l_{z3}\, l_\beta - l_s\, l_{z3}^2\, l_\beta - 3 l_t\, l_{z3}^2\, l_\beta - l_x\, l_{z3}^2\, l_\beta + \frac{11}{4} l_s^2\, l_{z4}\, l_\beta + 7 l_s\, l_t\, l_{z4}\, l_\beta +
\end{aligned}
$$

$$l_t^2\, l_{z4}\, l_\beta + 3\, l_s\, l_T\, l_{z4}\, l_\beta + 2\, l_t\, l_T\, l_{z4}\, l_\beta + \frac{1}{2}\, l_s\, l_x\, l_{z4}\, l_\beta + 3\, l_t\, l_x\, l_{z4}\, l_\beta - l_T\, l_x\, l_{z4}\, l_\beta -$$

$$\frac{1}{4}\, l_x^2\, l_{z4}\, l_\beta - 3\, l_s\, l_{z3}\, l_{z4}\, l_\beta - 6\, l_t\, l_{z3}\, l_{z4}\, l_\beta - 4\, l_T\, l_{z3}\, l_{z4}\, l_\beta + l_x\, l_{z3}\, l_{z4}\, l_\beta + 2\, l_{z3}^2\, l_{z4}\, l_\beta -$$

$$5\, l_s\, l_{z4}^2\, l_\beta - l_t\, l_{z4}^2\, l_\beta - 2\, l_x\, l_{z4}^2\, l_\beta + l_{z3}\, l_{z4}^2\, l_\beta + \frac{7}{3} l_{z4}^3\, l_\beta - \frac{7}{8} l_s^2\, l_\beta^2 - \frac{5}{2} l_s\, l_t\, l_\beta^2 - 2\, l_t^2\, l_\beta^2 -$$

$$\frac{5}{2}\, l_s\, l_T\, l_\beta^2 - 3\, l_t\, l_T\, l_\beta^2 + \frac{3}{4}\, l_s\, l_x\, l_\beta^2 - \frac{5}{2}\, l_t\, l_x\, l_\beta^2 - \frac{1}{2}\, l_T\, l_x\, l_\beta^2 - \frac{3}{8}\, l_x^2\, l_\beta^2 + \frac{5}{2}\, l_s\, l_{z3}\, l_\beta^2 +$$

$$3\, l_t\, l_{z3}\, l_\beta^2 + 2\, l_T\, l_{z3}\, l_\beta^2 - \frac{1}{2}\, l_x\, l_{z3}\, l_\beta^2 - l_{z3}^2\, l_\beta^2 + \frac{5}{2}\, l_s\, l_{z4}\, l_\beta^2 + 3\, l_t\, l_{z4}\, l_\beta^2 + 2\, l_T\, l_{z4}\, l_\beta^2 +$$

$$\frac{1}{2}\, l_x\, l_{z4}\, l_\beta^2 - 2\, l_{z3}\, l_{z4}\, l_\beta^2 - \frac{3}{2}\, l_{z4}^2\, l_\beta^2 - \frac{1}{3}\, l_s\, l_\beta^3 - \frac{2}{3}\, l_t\, l_\beta^3 - l_T\, l_\beta^3 - \frac{1}{3}\, l_x\, l_\beta^3 + l_{z3}\, l_\beta^3 +$$

$$l_{z4} l_\beta^3 + \frac{l_\beta^4}{6} + \Bigg(\frac{19}{8} l_x^2 - \frac{l_s^2}{8} - 9 l_t^2 + 5 l_T l_x + l_T l_{z3} + \frac{11}{2} l_x l_{z3} - l_{z3}^2 - 6 l_T l_{z4} + 2 l_x l_{z4} -$$

$$3\, l_{z3}\, l_{z4} - 7\, l_{z4}^2 + 2\, l_t\, \left(3\, l_T + 7\, l_{z3} + 5\, l_{z4} - 12\, l_\beta\right) + 6\, l_T\, l_\beta - 10\, l_x\, l_\beta - 2\, l_{z3}\, l_\beta +$$

$$14\, l_{z4}\, l_\beta - 13\, l_\beta^2 - \frac{1}{4}\, l_s\, \left(64\, l_t - 8\, l_T + 35\, l_x + 18\, l_{z3} - 44\, l_{z4} + 32\, l_\beta\right)\Bigg)\, \zeta(2) +$$

$$\left(-2\, l_s - 4\, l_t - l_T + l_x + 7\, l_{z3} + l_{z4}\right)\, \zeta(3) - \frac{35}{4}\zeta(4) - 2\, \mathrm{Li}_2^2\left(\frac{m^2}{z_5}\right) +$$

$$2\, \mathrm{Li}_2^2\left(\frac{-t(1-\beta)}{2\, m^2}\right) + \mathrm{Li}_2(x)\, \Bigg(2\, \mathrm{Li}_2\left(\frac{m^2}{z_5}\right) + 2\, \mathrm{Li}_2\left(\frac{-t(1-\beta)}{2\, m^2}\right) + \frac{l_s^2}{4} -$$

$$2\, l_t^2 + l_T^2 + l_T\, l_x + \frac{l_x^2}{4} + 2\, l_T\, l_{z3} + 3\, l_x\, l_{z3} + l_t\, \left(-4\, l_T - 3\, l_x + 2\, l_{z3}\right) +$$

$$l_s\, \left(l_t - l_T - \frac{9}{2} l_x - l_{z3} - l_{z4}\right) + 5\, l_x\, l_{z4} + l_{z4}^2 - 4\, l_x\, l_\beta - 6\, \zeta(2)\Bigg) + \mathrm{Li}_2\left(\frac{z_3}{z_4}\right) \times$$

$$\Bigg(-2\mathrm{Li}_2\left(\frac{m^2}{z_5}\right) - \frac{l_s^2}{4} + 4 l_t^2 + l_T\, l_x + \frac{3}{4} l_x^2 - \frac{l_s}{2}\left(6 l_t + 2 l_T + 3 l_x - 4 l_{z3} - 4 l_{z4}\right) +$$

$$2\, l_x\, l_{z4} - l_{z4}^2 - l_t\, \left(l_x - 2\, l_T + 2\, l_{z3} + 2\, l_{z4}\right) - 2\, l_x\, l_\beta - 6\, \zeta(2)\Bigg) +$$

$$\mathrm{Li}_2\left(\frac{-t(1-\beta)}{2\, m^2}\right)\, \Bigg(-2\mathrm{Li}_2\left(\frac{z_3}{z_4}\right) + \frac{1}{4}\Big(l_s^2 - 12 l_t^2 - 2 l_T^2 - 6 l_T\, l_x + l_x^2 - 12 l_T\, l_{z3} +$$

$$8\, l_x\, l_{z3} - 4\, l_x\, l_{z4} - 4\, l_{z4}^2 + l_s\, \left(-4\, l_t + 6\, l_T + 2\, l_x - 8\, l_{z3} + 4\, l_{z4} - 4\, l_\beta\right) +$$

$$4\, l_t\, \left(l_x + 4\, l_{z3} + 2\, l_{z4} - 2\, l_\beta\right) + 12\, l_x\, l_\beta + 8\, l_{z4}\, l_\beta - 4\, l_\beta^2\Big) + 12\, \zeta(2)\Bigg) +$$

$$\frac{1}{8}\, \mathrm{Li}_2\left(\frac{m^2 x}{-T}\right)\Big(-l_s^2 + 16 l_t l_T - l_x\left(4 l_T + l_x - 4 l_{z4}\right) - 2 l_s\left(2 l_T + l_x - 2 l_{z4}\right)\Big) + \frac{1}{4} \times$$

$$\mathrm{Li}_2(\frac{m^2}{z_5})\Big(3 l_s^2 + 4 l_t^2 + 2 l_T^2 + 2 l_T l_x + 7 l_x^2 + 8 l_T l_{z3} - 2 l_s (8 l_t + 3 l_T - l_x + 4 l_{z3} - 6 l_{z4})$$

$$+ 4 l_T l_{z4} + 4 l_x l_{z4} - 12 l_{z4}^2 + 8 l_t (l_{z3} + l_{z4} - l_\beta) + 8 l_x l_\beta - 8 l_{z3} l_\beta + 8 l_{z4} l_\beta - 4 l_\beta^2\Big) +$$

$$\frac{1}{8}\,\mathrm{Li}_2\left(\frac{T}{z_3}\right)\Big(9\,l_s^2-16\,l_t^2+l_x\,(-4\,l_T+l_x+12\,l_{z3}+8\,l_{z4}-8\,l_\beta)-$$

$$2l_s(4l_t-2l_T+5l_x+6l_{z3}+4l_{z4}-4l_\beta)-8l_t(2l_T+l_x-2l_{z3}-2l_{z4}+2l_\beta))+$$

$$\frac{1}{2}\mathrm{Li}_2(\frac{T}{m^2})\Big(2l_s^2-2l_Tl_x-2l_Tl_{z3}+3l_xl_{z3}+5l_{z3}^2-2l_t(l_x+2l_{z3})+2l_Tl_{z4}+3l_xl_{z4}+$$

$$4l_{z3}l_{z4}-l_{z4}^2-2l_xl_\beta-4l_{z3}l_\beta+l_s(2l_t-3l_x-7l_{z3}-l_{z4}+2l_\beta)\Big)+\frac{1}{4}\mathrm{Li}_2(-x)\times$$

$$\Big(5l_s^2-16l_t^2+l_x^2-4l_Tl_{z3}+6l_xl_{z3}-6l_{z3}^2-4l_Tl_{z4}+2l_xl_{z4}-8l_{z3}l_{z4}+2l_{z4}^2-4l_x\,l_\beta+$$

$$8l_{z3}l_\beta+2l_s(-6l_t+2l_T-2l_x+l_{z3}-3l_{z4}+2l_\beta)-4l_t(l_x-6l_{z3}-4l_{z4}+4l_\beta))+$$

$$\mathrm{Li}_3\left(\frac{z_4}{T}\right)(-l_T+l_{z4})+2\left(\mathrm{Li}_3\left(\frac{z_3}{s\beta}\right)+\mathrm{Li}_3\left(\frac{z_5}{t\beta}\right)\right)(2\,l_t+2\,l_T-l_{z3}-l_{z4})+$$

$$2\mathrm{Li}_3\left(\frac{-1+\beta}{2\beta}\right)(2\,l_s-2\,l_t-2\,l_T-l_{z3}-l_{z4})+$$

$$\mathrm{Li}_3\left(\frac{m^2}{z_5}\right)\left(-\frac{17}{2}l_s-8\,l_t+4\,l_T-\frac{l_x}{2}+8\,l_{z3}+9\,l_{z4}-8\,l_\beta\right)+$$

$$\frac{1}{2}\,\mathrm{Li}_3\left(\frac{T}{z_6}\right)(-5\,l_s-4\,l_t-2\,l_T+l_x+6\,l_{z3}+4\,l_{z4}-4\,l_\beta)+$$

$$2\,\mathrm{Li}_3\left(\frac{z_3}{z_4}\right)(-l_s+l_t+l_T+l_x-l_\beta)+\mathrm{Li}_3\left(\frac{z_3}{t}\right)(-l_s-2\,l_T-3\,l_x+2\,l_{z3})+$$

$$\frac{1}{2}\mathrm{Li}_3\left(\frac{z_6}{m^2}\right)(l_s+8l_t-8l_T-l_x-6l_{z3}-4l_{z4})+\mathrm{Li}_3\left(\frac{z4}{t}\right)(l_s+4l_t+2l_T-3l_x-$$

$$2l_{z3}-4l_{z4})+\frac{1}{2}\mathrm{Li}_3\left(\frac{z_5}{T}\right)(l_s+2l_T+l_x-2l_{z4})+2\Big(\mathrm{Li}_3\left(\frac{-x^2}{1-x^2}\right)-$$

$$2\mathrm{Li}_3\left(\frac{1-\beta}{2}\right)-\mathrm{Li}_3\left(\frac{m^2(1-\beta)}{2\,z_5}\right)+\mathrm{Li}_3\left(\frac{-t(1-\beta)}{2\,z_5}\right)-2\mathrm{Li}_3\left(\frac{T}{m^2}\right)\Big)\,l_T+$$

$$2\left(2\mathrm{Li}_3\left(\frac{T}{m^2}\right)+\mathrm{Li}_3\left(\frac{2\,z_6}{m^2(1+\beta)}\right)-\mathrm{Li}_3\left(\frac{-2\,z_6}{t(1+\beta)}\right)+\mathrm{Li}_3\left(\frac{z_6}{z_5}\right)\right)(l_s+l_t+$$

$$l_T-l_{z3}-l_{z4}+l_\beta)-2\mathrm{Li}_3\left(\frac{z_6}{z_5}\right)l_x-\mathrm{Li}_3\left(\frac{T}{z_3}\right)l_{z3}+\left(\mathrm{Li}_3\left(\frac{T}{z_3}\right)+2\,\mathrm{Li}_3(-x)\right)\times$$

$$(2l_s+2l_t+l_T-2l_{z3}-2l_{z4}+2l_\beta)+2\mathrm{Li}_3(x)\,(4l_s+3l_t+l_T+l_x-2l_{z3}-6l_{z4}+$$

$$3l_\beta)+2\mathrm{Li}_4(x)-\mathrm{Li}_4\left(\frac{T}{z_3}\right)-4\mathrm{Li}_4\left(\frac{z_3}{t}\right)+4\mathrm{Li}_4\left(\frac{z_4}{t}\right)-\mathrm{Li}_4\left(\frac{z_4}{T}\right)+\mathrm{Li}_4\left(\frac{z_5}{T}\right)+$$

$$2\mathrm{Li}_4\left(\frac{s(1-\beta)}{-2t}\right)+3\mathrm{Li}_4\left(\frac{s(1-\beta)}{2\,z_4}\right)+4\mathrm{Li}_4\left(\frac{-1+\beta}{2\,\beta}\right)+2\mathrm{Li}_4\left(\frac{-2t}{s(1+\beta)}\right)+$$

$$4\mathrm{Li}_4\left(\frac{2\,\beta}{1+\beta}\right)+\mathrm{Li}_4\left(\frac{sT(1+\beta)}{2\,m^2\,z_3}\right)+3\mathrm{Li}_4\left(\frac{2\,z_3}{s(1+\beta)}\right)+$$

$$\frac{1}{2}L_{-++}\left(1,\frac{m^2}{-T},\frac{m^2}{-T},\frac{s(1-\beta)}{-2z_4}\right)-\frac{1}{2}L_{-++}\left(1,\frac{m^2}{-T},\frac{m^2}{-T},\frac{s(1+\beta)}{-2z_3}\right)+$$

$$2L_{-++}\left(1,\frac{m^2}{-T},\frac{s(1-\beta)}{-2z_4},0\right)+L_{-++}\left(1,-\frac{t}{z_3},-\frac{t}{z_3},-\frac{t}{T}\right)-$$

$$L_{-++}\left(1,\frac{s(1-\beta)}{-2z_4},\frac{m^2}{-T},\frac{s(1-\beta)}{-2z_4}\right)-L_{-++}\left(1,\frac{s(1-\beta)}{-2z_4},\frac{m^2}{-T},\frac{s(1+\beta)}{-2z_3}\right)+$$

$$L_{-++}\left(1,\frac{s(1-\beta)}{-2z_4},\frac{s(1-\beta)}{-2z_4},0\right)-\frac{1}{2}L_{-++}\left(1,\frac{s(1-\beta)}{-2z_4},\frac{s(1-\beta)}{-2z_4},\frac{s(1+\beta)}{-2z_3}\right)+$$

$$\frac{1}{2}L_{-++}\left(\frac{t}{T},0,0,-\frac{t}{z_3}\right)-\frac{1}{2}L_{-++}\left(\frac{t}{T},0,0,-\frac{t}{z_4}\right)+2L_{-++}\left(\frac{t}{T},0,-\frac{t}{z_3},-1\right)-$$

$$L_{-++}\left(\frac{t}{T},0,-\frac{t}{z_3},-\frac{t}{z_3}\right)-L_{-++}\left(\frac{t}{T},0,-\frac{t}{z_3},-\frac{t}{z_4}\right)-L_{-++}\left(\frac{t}{T},-\frac{t}{z_3},-\frac{t}{z_3},-\frac{t}{z_4}\right)+$$

$$3L_{-++}(\frac{t}{z_4},0,0,-1)-\frac{3}{2}L_{-++}\left(\frac{t}{z_4},0,0,-\frac{t}{z_3}\right)-2L_{-++}\left(\frac{t}{z_4},0,-\frac{t}{z_3},-1\right)+$$

$$L_{-++}\left(\frac{t}{z_4},0,-\frac{t}{z_3},-\frac{t}{z_3}\right)+3L_{-++}\left(\frac{t}{z_4},0,-\frac{t}{z_3},-\frac{t}{z_4}\right)+$$

$$\frac{1}{2}L_{-++}\left(\frac{t}{z_4},-\frac{t}{z_3},-\frac{t}{z_3},-\frac{t}{T}\right)+2L_{-++}\left(\frac{s(1+\beta)}{2\,z_3},\frac{m^2}{-T},0,\frac{m^2}{-T}\right)-$$

$$2L_{-++}\left(\frac{s(1+\beta)}{2\,z_3},\frac{m^2}{-T},0,\frac{s(1-\beta)}{-2z_4}\right)-2L_{-++}\left(\frac{s(1+\beta)}{2\,z_3},\frac{m^2}{-T},0,\frac{s(1+\beta)}{-2z_3}\right)+$$

$$\frac{1}{2}L_{-++}\left(\frac{s(1+\beta)}{2\,z_3},\frac{m^2}{-T},\frac{m^2}{-T},\frac{s(1-\beta)}{-2z_4}\right)+2L_{-++}\left(\frac{s(1+\beta)}{2\,z_3},\frac{s(1-\beta)}{-2z_4},0,\frac{m^2}{-T}\right)-$$

$$\frac{1}{2}L_{-++}\left(\frac{s(1+\beta)}{2\,z_3},\frac{s(1-\beta)}{-2z_4},\frac{s(1-\beta)}{-2z_4},\frac{m^2}{-T}\right)-4L_{+}\left(0,\frac{z_3}{s\beta},-\frac{z_3z_4}{st\beta},-1\right)+$$

$$4L_{+}\left(0,\frac{z_3(1-\beta)}{-2t\beta},\frac{z_3z_4}{st\beta},\frac{m^2}{-T}\right)+2L_{+}\left(\frac{m^2}{-T},\frac{z_3(1-\beta)}{-2t\beta},\frac{z_3z_4}{st\beta},\frac{s(1-\beta)}{-2z_4}\right)+$$

$$2L_{+}\left(-\frac{t}{z_3},0,\frac{z_4}{t},-\frac{t}{z_4}\right)+2L_{+}\left(-\frac{t}{z_3},0,\frac{z_3}{t},-\frac{t}{z_4}\right)+2L_{+}\left(-\frac{t}{z_3},\frac{z_3}{s\beta},-\frac{z_3z_4}{st\beta},-\frac{t}{T}\right)+$$

$$\frac{3}{2}L_{+++}\left(0,0,-\frac{t}{z_3},-\frac{t}{z_4}\right)-2L_{+++}\left(0,\frac{m^2}{-T},\frac{s(1-\beta)}{-2z_4},\frac{m^2}{-T}\right)+$$

$$2L_{+++}\left(0,\frac{m^2}{-T},\frac{s(1-\beta)}{-2z_4},\frac{s(1-\beta)}{-2z_4}\right)+2L_{+++}\left(0,\frac{m^2}{-T},\frac{s(1-\beta)}{-2z_4},\frac{s(1+\beta)}{-2z_3}\right)+$$

$$L_{+++}\left(0,-\frac{t}{z_3},-\frac{t}{z_3},-1\right)-L_{+++}\left(\frac{m^2}{-T},\frac{m^2}{-T},0,\frac{s(1-\beta)}{-2z_4}\right)+$$

$$L_{+++}\left(\frac{m^2}{-T},\frac{m^2}{-T},0,\frac{s(1+\beta)}{-2z_3}\right)-\frac{1}{2}L_{+++}\left(\frac{m^2}{-T},\frac{m^2}{-T},\frac{s(1-\beta)}{-2z_4},\frac{s(1+\beta)}{-2z_3}\right)-$$

$$3L_{+++}\left(-\frac{t}{z_3},0,0,-1\right)+\frac{3}{2}L_{+++}\left(-\frac{t}{z_3},-\frac{t}{z_3},0,-\frac{t}{z_4}\right)+$$

$$L_{+++}\left(\frac{s(1-\beta)}{-2z_4},\frac{s(1-\beta)}{-2z_4},0,\frac{m^2}{-T}\right)\Bigg];$$

$$\begin{aligned}
\operatorname{Im} D_1^{(-1)} &= \pi/(st\beta), \qquad\qquad \operatorname{Im} D_1^{(0)} = -2\pi\ln(-t\beta/m^2)/(st\beta), \qquad (4.52)\\
\operatorname{Im} D_1^{(1)} &= \frac{\pi}{st\beta}\left[-\frac{3}{4}\,l_s^2+2\,l_t^2-\frac{3}{4}\,l_x^2-l_xl_{z3}+l_xl_{z4}-l_{z4}^2+l_s\,(l_t+l_x/2+l_{z3}+l_{z4})-\right.\\
&\left. l_t\,(l_x+2\,l_{z3}-4\,l_\beta)+2\,l_\beta^2-2\zeta(2)-2\mathrm{Li}_2\left(\frac{m^2}{z_5}\right)-2\mathrm{Li}_2\left(\frac{t(-1+\beta)}{2m^2}\right)\right],\\
\operatorname{Im} D_1^{(2)} &= \frac{\pi}{st\beta}\left[\frac{1}{12}\left(7\,l_s^3-l_x^3-3\,l_s^2\,(3\,l_x+4\,l_{z3}+4\,l_{z4}-6\,l_\beta)+6\,l_x^2\,(-2\,l_{z3}+3\,l_\beta)+\right.\right.\\
&12\,l_x\left(l_{z4}^2-2\,(-l_{z3}+l_{z4})\,l_\beta+2\,l_t\,(-l_{z3}+l_\beta)\right)-\\
&8\,l_\beta\left(6\,l_t^2-3\,l_{z4}^2+2\,l_\beta^2+6\,l_t\,(-l_{z3}+l_\beta)\right)-3\,l_s\left(8\,l_t^2-5\,l_x^2-4\,l_{z4}^2-\right.\\
&\left.\left.8\,l_t\,(l_x+l_{z3}-l_\beta)+8\,l_{z3}\,l_\beta+8\,l_{z4}\,l_\beta+4\,l_x\,(-2\,l_{z3}+2\,l_{z4}+l_\beta)\right)\right)+\\
&4\,(l_t+l_x+l_\beta)\,\zeta(2)+2\,\mathrm{Li}_2(x)\,l_x+2\,\mathrm{Li}_2\left(\frac{z3}{z4}\right)\,l_x+\\
&2\,\mathrm{Li}_2\left(\frac{t(-1+\beta)}{2\,m^2}\right)(l_s-l_x+2\,l_\beta)+2\,\mathrm{Li}_2\left(\frac{m^2}{z_5}\right)(l_s+l_x+2\,l_\beta)-\\
&\left.2\,\mathrm{Li}_3(x)-4\,\mathrm{Li}_3\left(\frac{z_3}{t}\right)+2\,\mathrm{Li}_3\left(\frac{z_3}{z_4}\right)-4\,\mathrm{Li}_3\left(\frac{z_4}{t}\right)-2\,\mathrm{Li}_3\left(\frac{z_6}{z_5}\right)\right].
\end{aligned}$$



FIG. 3. Massive box D_2 with one massive propagator.

Next we turn to the second four-point function D_2 with one massive propagator shown in Fig. 3 which is defined by

$$D_2 \equiv D(-p_2,p_4,p_3,0,0,m,0).$$

We substitute the appropriate values of momenta and masses for the D_2 integral into the general kernel expression (4.40) and obtain

$$K = ac\tilde{t}-bd\tilde{s}+c^2-i\delta. \tag{4.53}$$

In order to simplify the first integration over the Feynman parameter x_3 we choose $\{a, b, c, d\}$ as $\{x_1x_2(1-x_3),\, x_1x_2x_3,\, x_1(1-x_2),\, 1-x_1\}$. After x_3-integration we write the result for the integrand as a sum of two parts $I^{D_2}_{x_1x_2} + II^{D_2}_{x_1x_2}$:

$$I^{D_2}_{x_1x_2} = -\frac{x_1\left[x_1^2(1-x_2)(1+(\tilde{t}-1)x_2) - i\delta\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{s}(1-x_1)+\tilde{t}x_1(1-x_2)]}, \tag{4.54}$$

$$II^{D_2}_{x_1x_2} = \frac{x_1\left[-\tilde{s}x_1x_2 + x_1^2(1+(\tilde{s}-2)x_2+x_2^2) - i\delta\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{s}(1-x_1)+\tilde{t}x_1(1-x_2)]}, \tag{4.55}$$

where again the two terms derive from the indefinite integral (or primitive) evaluated at the upper and lower boundary of x_3. First consider the integration of $I^{D_2}_{x_1x_2}$. One notes that its numerator is not negative on the integration path since $\tilde{t} > 1$, which implies there is no imaginary contribution coming from $I^{D_2}_{x_1x_2}$. Therefore, we omit the $i\delta$ term for the remaining integration. After integration over x_1 we arrive at

$$I^{D_2}_{x_2} = \frac{\left[(1-x_2)\left(1+(\tilde{t}-1)\,x_2\right)\right]^{-1-\varepsilon}\ {}_2F_1(1,-2\varepsilon,1-2\varepsilon,1-A)}{2\tilde{s}\,(1+\varepsilon)\,\varepsilon}, \tag{4.56}$$

where we have defined $A \equiv \tilde{t}\,(1-x_2)/\tilde{s}$ and ${}_2F_1$ is a hypergeometric function. The above expression is singular at the upper integration limit $x_2 = 1$. In order to regularize this singularity we have to find a suitable subtraction term.

First note that the ε-expansion of the hypergeometric function reads

$$\begin{aligned} F_1(1,-2\varepsilon,1-2\varepsilon,1-A) = {} & 1 + 2\varepsilon\,\ln A - 4\varepsilon^2\,\mathrm{Li}_2(1-A) - 8\varepsilon^3\,\mathrm{Li}_3(1-A) \\ & -16\varepsilon^4\,\mathrm{Li}_4(1-A) + \mathcal{O}(\varepsilon^5). \end{aligned} \tag{4.57}$$

To obtain a suitable subtraction term we substitute (4.57) into (4.56) and replace x_2 by 1 everywhere in $I^{D_2}_{x_2}$ except for terms that diverge. Therefore, our subtraction term can be defined as

$$I^{D_2,s}_{x_2} = \frac{(1-x_2)^{-1-\varepsilon}\,\tilde{t}^{-1-\varepsilon}\,[1+2\varepsilon\,\ln A - 4\varepsilon^2\zeta(2) - 8\varepsilon^3\,\zeta(3) - 16\varepsilon^4\zeta(4)]}{2\tilde{s}\,(1+\varepsilon)\,\varepsilon}. \tag{4.58}$$

Our subtraction term is simple enough to be integrated analytically over x_2 giving the result

$$I^{D_2,s} = \frac{\tilde{t}^{-1-\varepsilon}\left(-3 - 2\varepsilon\,\ln\frac{\tilde{t}}{\tilde{s}} + 4\varepsilon^2\,\zeta(2) + 8\varepsilon^3\,\zeta(3) + 16\varepsilon^4\,\zeta(4)\right)}{2\,(1+\varepsilon)\,\varepsilon^2\,\tilde{s}} \tag{4.59}$$

which can readily be expanded in ε.

Next we turn to the remaining finite integral $I^{D_2}_{x_2} - I^{D_2,s}_{x_2}$. We expand $I^{D_2}_{x_2} - I^{D_2,s}_{x_2}$ up to second order in ε and integrate over x_2 after the expansion.

Next consider the second integrand $II^{D_2}_{x_1x_2}$ (4.55). The term in the numerator in square brackets raised to the power $(-1-\varepsilon)$ changes sign on the integration path. It means that the corresponding integral has an imaginary contribution. We can rewrite $II^{D_2}_{x_1x_2}$ as follows:

$$II^{D_2}_{x_1x_2} = \frac{x_1^{-\varepsilon}(\tilde{s}x_2)^{-1-\varepsilon}\left(-1-i\delta + x_1(\frac{(1-x_2)^2}{\tilde{s}x_2}+1)\right)^{-1-\varepsilon}}{(1+\varepsilon)\tilde{s}\left(1-x_1(1-\frac{\tilde{t}}{\tilde{s}}(1-x_2))\right)}, \tag{4.60}$$

The integration of $II^{D_2}_{x_1x_2}$ over x_1 is more difficult because of the additional term $x_1^{-\varepsilon}$. We proceed by expanding $x_1^{-\varepsilon}$ as $(1-\varepsilon\ln x_1 + \frac{\varepsilon^2}{2}\ln^2 x_1 + ...)$. One can see that only the first term of this expansion gives rise to a divergence in the subsequent integration. As we will need to find a subtraction term for this term we will treat it separately. For the remaining terms we do an overall ε-expansion of (4.60) and then perform the remaining integrations. We therefore substitute $x_1^{-\varepsilon}$ by 1. The integral looks simpler (we denote it by $II^0_{x_1x_2}$). Integration over x_1 yields

$$\begin{aligned} II^0_{x_2} = {} & \frac{(\tilde{s}x_2)^{-1-\varepsilon}x_2(-1-i\delta)^{-\varepsilon}\,{}_2F_1(1,-\varepsilon,1-\varepsilon,-\frac{x_2(\tilde{s}+\tilde{t}(-1+x_2))}{(1-x_2)(1+(\tilde{t}-1)x_2)})}{\varepsilon(1+\varepsilon)(1-x_2)(1+(\tilde{t}-1)x_2)} \\ & -\frac{(\tilde{s}x_2)^{-1-\varepsilon}x_2\,(1-x_2)^{-2\varepsilon}\,{}_2F_1(1,-\varepsilon,1-\varepsilon,\frac{(1-x_2)(\tilde{s}+\tilde{t}(-1+x_2))}{\tilde{s}(1+(-1+\tilde{t})x_2)})}{\varepsilon(1+\varepsilon)(1-x_2)(1+(\tilde{t}-1)x_2)(\tilde{s}x_2)^{-\varepsilon}}. \end{aligned} \tag{4.61}$$

Note that we omit the imaginary shifts $i\delta$ in the arguments of the hypergeometric functions ${}_2F_1$, as the branch cuts of ${}_2F_1$ are never crossed in the physical region. If we would directly integrate the above expression we would have a divergence at $x_2 \to 1$. We must therefore define a subtraction term. If one uses $II^0_{x_2}$ in the present form the definition is rather difficult: as $x_2 \to 1$ the argument of the first function ${}_2F_1$ goes to infinity. To circumvent this problem we can use one of the relations between hypergeometric functions to transform the argument of the function. As a result we pull out the divergent term as an overall factor multiplying the hypergeometric function with a transformed argument. The whole expression can be rewritten as

$$\begin{aligned} II^0_{x_2} = {} & -\frac{(1-x_2)^{-1-2\varepsilon}\,{}_2F_1(1,-\varepsilon,1-\varepsilon,\frac{\left(\tilde{s}+\tilde{t}\,(-1+x_2)\right)(1-x_2)}{\tilde{s}\left(1+\left(-1+\tilde{t}\right)x_2\right)})}{\tilde{s}\,\left(1+\left(-1+\tilde{t}\right)\,x_2\right)\,(1+\varepsilon)\,\varepsilon} + \\ & \frac{(\tilde{s}(1-x_2))^{-1-\varepsilon}\,x_2^{-\varepsilon}\left(1+\left(-1+\tilde{t}\right)x_2\right)^{-1-\varepsilon}(-1-i\,\delta)^{-\varepsilon}\,{}_2F_1(-\varepsilon,-\varepsilon,1-\varepsilon,\frac{\left(\tilde{s}-\tilde{t}\,(1-x_2)\right)x_2}{(1-x_2)^2+\tilde{s}\,x_2})}{\left((1-x_2)^2+\tilde{s}\,x_2\right)^{-\varepsilon}(1+\varepsilon)\,\varepsilon}. \end{aligned} \tag{4.62}$$

Now all the poles arise from the factors $(1-x_2)^{-1-2\varepsilon}$ and $(1-x_2)^{-1-\varepsilon}$, and we can derive the necessary subtraction term following the above procedure. We briefly mention that when $x_2 = 1$ the first hypergeometric function (in the first line of Eq. (4.62)) takes the value ${}_2F_1 = 1$. The second hypergeometric function takes the value $-\varepsilon\pi/\sin(-\varepsilon\pi)$ which, in turn, can be expanded to order ε^4 as $1+\varepsilon^2\zeta(2)+7\varepsilon^4\zeta(4)/4$. Thus, the subtraction term reads

$$II^{0,s}_{x_2} = -\frac{(1-x_2)^{-1-2\varepsilon}}{(1+\varepsilon)\varepsilon\tilde{s}\tilde{t}} + \frac{\left(1+\varepsilon^2\zeta(2)+\frac{7\varepsilon^4\zeta(4)}{4}\right)\tilde{t}^{-1-\varepsilon}\,(1-x_2)^{-1-\varepsilon}\,(-1-i\delta)^{-\varepsilon}}{(1+\varepsilon)\,\varepsilon\,\tilde{s}}. \qquad (4.63)$$

Integrating this subtraction term we arrive at

$$II^{0,s} = \frac{1}{2(1+\varepsilon)\varepsilon^2\tilde{s}\tilde{t}} - \frac{\left(1+\varepsilon^2\zeta(2)+\frac{7\varepsilon^4\zeta(4)}{4}\right)\tilde{t}^{-1-\varepsilon}\,(-1-i\delta)^{-\varepsilon}}{(1+\varepsilon)\varepsilon^2\tilde{s}}, \qquad (4.64)$$

which can finally be expanded up to ε^2.

Now that we have found a suitable subtraction term we can proceed with the remaining terms. We subtract from $II^0_{x_2}$ (Eq. (4.62)) the subtraction term $II^{0,s}_{x_2}$ (Eq. (4.63)). Since the result is convergent with regard to the integration over x_2, we can expand the result in terms of ε before integration which greatly simplifies the problem. We then do the last integration. The difference $II^0_{x_2} - II^{0,s}_{x_2}$ must be expanded up to third order in ε. The reason for this is that we have already one pole $\sim 1/\varepsilon$ after the x_1-integration. Therefore, in order to get results up to second order the hypergeometric functions have to be expanded to third order. The expansion for one of the hypergeometric functions is done using (4.57). For the ε-expansion of the second hypergeometric function one gets

$$\begin{aligned}
&{}_2F_1(-\varepsilon,-\varepsilon,1-\varepsilon,z) = 1+\varepsilon^2\,\mathrm{Li}_2(z) && (4.65)\\
&-\varepsilon^3\left(\frac{1}{2}\ln^2(1-z)\ln z + \ln(1-z)\mathrm{Li}_2(1-z) - \mathrm{Li}_3(1-z) - Li_3(z) + \zeta(3)\right).
\end{aligned}$$

Using these results for the ε-expansions we expand $II^0_{x_2} - II^{0,s}_{x_2}$ up to ε^2 and integrate the resulting expression. Finally, carefully collecting all the relevant pieces, we arrive at the final result for our second four-point function. In order to reduce the length of the final result for D_2 we introduce four more abbreviations. We write

$$D \equiv m^2 s - tu, \qquad l_D \equiv \ln\frac{-D}{m^4}, \qquad l_u \equiv \ln\frac{-u}{m^2}, \qquad l_U \equiv \ln\frac{-U}{m^2} \qquad (4.66)$$

Our result for the four-point box diagram D_2 reads:

$$\mathrm{Re}\,D_2^{(-2)} = 2/(st), \qquad \mathrm{Re}\,D_2^{(-1)} = -[l_s+2l_t]/(st), \qquad (4.67)$$

$$\begin{aligned}
\mathrm{Re}\,D_2^{(0)} &= [2\,l_s\,l_t - 5\,\zeta(2)]/(st),\\
\mathrm{Re}\,D_2^{(1)} &= \frac{1}{s\,t}\left[\frac{1}{2}\,l_s^3 + 3\,l_t^3 + \frac{l_x^3}{12} - l_t^2\,(4\,l_T + 3\,l_x + 4\,l_{z3} - l_{z4}) + \frac{1}{2}\,l_x^2\,l_{z4} - l_x\,l_{z4}^2 + \frac{2}{3}\,l_{z4}^3 - \right.\\
&\quad \frac{1}{4}l_s^2\,(6l_t + 3l_x + 4l_{z3} + 2l_{z4}) - l_s\left(2l_t^2 - \frac{1}{2}l_x^2 - l_x l_{z3} - 2l_t\,(l_x + 2l_{z3} + l_{z4})\right) -
\end{aligned}$$

$$
\begin{aligned}
& l_t \left(l_T^2 + \frac{1}{2} l_x^2 + 2 l_x l_{z3} - 2 l_x l_{z4} - 2 l_T \left(l_x + l_{z3} + l_{z4}\right) + l_{z3}^2 + 3 l_{z4}^2\right) - (l_s - 2 l_t + \\
& l_x - 4 l_{z4})\, \zeta(2) + \zeta(3) + 2 \left(\mathrm{Li}_2\left(\frac{m^2}{z_5}\right) + \mathrm{Li}_2\left(\frac{-t(1-\beta)}{2\, m^2}\right)\right) (l_s - 2\, l_t) - \\
& 4 \mathrm{Li}_2\left(\frac{T}{m^2}\right) l_t - 2 \mathrm{Li}_2\left(\frac{m^2 x}{-T}\right) l_t - 2 \mathrm{Li}_2\left(\frac{T}{z_3}\right) l_t - 4 \mathrm{Li}_3\left(\frac{m^2}{-t}\right) + 2 \mathrm{Li}_3(-x) - \\
& 2 \mathrm{Li}_3\left(\frac{z_3}{t}\right) - 2 \mathrm{Li}_3\left(\frac{z_4}{t}\right) - 4 \mathrm{Li}_3\left(\frac{m^2}{z_5}\right) - 4 \mathrm{Li}_3\left(\frac{z_6}{m^2}\right) + 2 \mathrm{Li}_3\left(\frac{m^2(1-\beta)}{2\, z_5}\right) - \\
& 2 \mathrm{Li}_3\left(\frac{-t(1-\beta)}{2\, z_5}\right) + 2 \mathrm{Li}_3\left(\frac{2\, z_6}{m^2(1+\beta)}\right) - 2 \mathrm{Li}_3\left(\frac{2\, z_6}{-t(1+\beta)}\right)\Biggr], \\
\mathrm{Re}\, D_2^{(2)} \quad = \quad & \frac{1}{s\, t} \Biggl[-\frac{l_s^4}{48} - \frac{17}{24} l_s^3 l_t - \frac{9}{2} l_s^2 l_t^2 - 3 l_s l_t^3 + \frac{13}{6} l_s^3 l_T + \frac{5}{2} l_s^2 l_t l_T + \frac{3}{2} l_t^3 l_T - \frac{1}{2} l_s^2 l_T^2 - \frac{7}{2} l_t^2 l_T^2 \\
& -3 l_t l_T^3 - \frac{25}{24} l_t^4 - \frac{3}{4} l_s^3 l_u - \frac{13}{4} l_s^2 l_t l_u + 7 l_s l_t^2 l_u - 3 l_t^3 l_u - \frac{5}{4} l_s^2 l_T l_u + 2 l_s l_t l_T l_u - \\
& 2 l_t^2 l_T l_u + \frac{1}{2} l_s^2 l_u^2 - \frac{3}{2} l_s l_t l_u^2 + l_t^2 l_u^2 - \frac{1}{2} l_s l_T l_u^2 + l_t l_T l_u^2 + \frac{1}{3} l_s l_u^3 - \frac{1}{3} l_t l_u^3 - \frac{1}{4} l_s^3 l_x - \\
& \frac{23}{8} l_s^2 l_t l_x + 3 l_s l_t^2 l_x - \frac{7}{3} l_t^3 l_x - \frac{7}{4} l_s^2 l_T l_x - 3 l_s l_t l_T l_x - 3 l_t^2 l_T l_x + 3 l_t l_T^2 l_x - \frac{1}{2} l_s^2 l_u l_x + \\
& \frac{1}{2} l_s l_t l_u l_x + \frac{5}{2} l_s l_T l_u l_x + \frac{1}{2} l_s l_u^2 l_x - \frac{1}{2} l_t l_u^2 l_x - \frac{1}{2} l_T l_u^2 l_x + \frac{1}{4} l_s^2 l_x^2 + \frac{19}{8} l_s l_t l_x^2 + \frac{3}{2} l_t^2 l_x^2 - \\
& 4 l_s l_T l_x^2 - \frac{9}{2} l_t l_T l_x^2 - \frac{1}{2} l_T^2 l_x^2 - \frac{3}{4} l_s l_u l_x^2 + \frac{3}{4} l_t l_u l_x^2 - \frac{1}{4} l_T l_u l_x^2 + \frac{5}{6} l_s l_x^3 + \frac{53}{24} l_t l_x^3 - \\
& \frac{37}{12} l_T l_x^3 - \frac{13}{16} l_x^4 - \frac{29}{24} l_s^3 l_{z3} + \frac{19}{4} l_s^2 l_t l_{z3} + 11 l_s l_t^2 l_{z3} + \frac{10}{3} l_t^3 l_{z3} - \frac{13}{4} l_s^2 l_T l_{z3} - \\
& 10 l_s l_t l_T l_{z3} - 9 l_t^2 l_T l_{z3} + 7 l_t l_T^2 l_{z3} + 2 l_s^2 l_u l_{z3} - 2 l_s l_t l_u l_{z3} + \frac{5}{8} l_s^2 l_x l_{z3} + \frac{11}{2} l_s l_t l_x l_{z3} + \\
& l_t^2 l_x l_{z3} - \frac{5}{2} l_s l_T l_x l_{z3} - 10 l_t l_T l_x l_{z3} + \frac{19}{8} l_s l_x^2 l_{z3} + \frac{7}{4} l_t l_x^2 l_{z3} - \frac{5}{4} l_T l_x^2 l_{z3} - \frac{19}{24} l_x^3 l_{z3} - \\
& \frac{5}{2} l_s l_t l_{z3}^2 - \frac{5}{2} l_t^2 l_{z3}^2 + \frac{7}{2} l_s l_T l_{z3}^2 + l_t l_T l_{z3}^2 - l_s l_u l_{z3}^2 + l_t l_u l_{z3}^2 - \frac{1}{2} l_s l_x l_{z3}^2 + \frac{7}{2} l_t l_x l_{z3}^2 + \\
& \frac{3}{2} l_T l_x l_{z3}^2 - \frac{1}{2} l_x^2 l_{z3}^2 + \frac{1}{3} l_s l_{z3}^3 + l_t l_{z3}^3 - l_T l_{z3}^3 - \frac{2}{3} l_x l_{z3}^3 - l_s^3 l_{z4} + 5 l_s^2 l_t l_{z4} + 7 l_s l_t^2 l_{z4} + \\
& \frac{16}{3} l_t^3 l_{z4} - \frac{5}{2} l_s^2 l_T l_{z4} - 4 l_s l_t l_T l_{z4} - 2 l_t^2 l_T l_{z4} + 8 l_t l_T^2 l_{z4} + 3 l_s^2 l_u l_{z4} - 3 l_s l_t l_u l_{z4} - \\
& l_s l_T l_u l_{z4} + l_s l_u^2 l_{z4} - l_t l_u^2 l_{z4} - l_T l_u^2 l_{z4} + 4 l_s^2 l_x l_{z4} - 3 l_s l_t l_x l_{z4} + 5 l_t^2 l_x l_{z4} + 4 l_t l_T l_x l_{z4} \\
& -l_s l_u l_x l_{z4} + l_t l_u l_x l_{z4} - 3 l_T l_u l_x l_{z4} + \frac{3}{2} l_s l_x^2 l_{z4} + l_t l_x^2 l_{z4} + \frac{5}{2} l_T l_x^2 l_{z4} + \frac{1}{6} l_x^3 l_{z4} + \\
& 2 l_s^2 l_{z3} l_{z4} - 14 l_s l_t l_{z3} l_{z4} - 4 l_t^2 l_{z3} l_{z4} + 6 l_s l_T l_{z3} l_{z4} + 6 l_t l_T l_{z3} l_{z4} - 2 l_s l_u l_{z3} l_{z4} + \\
& 2 l_t l_u l_{z3} l_{z4} - 2 l_s l_x l_{z3} l_{z4} - 2 l_t l_x l_{z3} l_{z4} + 2 l_T l_x l_{z3} l_{z4} - 4 l_x^2 l_{z3} l_{z4} + 3 l_t l_{z3}^2 l_{z4} -
\end{aligned}
$$

$$
\begin{aligned}
&2l_T l_{z3}^2 l_{z4} - l_s^2 l_{z4}^2 - 2l_s l_t l_{z4}^2 - \frac{9}{2} l_t^2 l_{z4}^2 + 2l_s l_T l_{z4}^2 - 3l_t l_T l_{z4}^2 - 2l_s l_u l_{z4}^2 + 2l_t l_u l_{z4}^2 + \\
&l_T l_u l_{z4}^2 - l_s l_x l_{z4}^2 - 2l_t l_x l_{z4}^2 + l_T l_x l_{z4}^2 - l_x^2 l_{z4}^2 + l_s l_{z3} l_{z4}^2 + 2l_t l_{z3} l_{z4}^2 - 2l_T l_{z3} l_{z4}^2 + \\
&l_x l_{z3} l_{z4}^2 + \frac{1}{3} l_s l_{z4}^3 + \frac{5}{3} l_t l_{z4}^3 - \frac{2}{3} l_T l_{z4}^3 - \frac{1}{3} l_x l_{z4}^3 - 2l_s^3 l_\beta - l_s^2 l_t l_\beta - l_s l_t^2 l_\beta - l_s^2 l_T l_\beta - \\
&\frac{3}{2} l_s^2 l_x l_\beta - l_t^2 l_x l_\beta - 2l_s l_T l_x l_\beta + l_s l_x^2 l_\beta + l_t l_x^2 l_\beta - l_T l_x^2 l_\beta - \frac{1}{2} l_x^3 l_\beta + 2l_s^2 l_{z3} l_\beta + \\
&2l_s l_t l_{z3} l_\beta + 2l_s l_x l_{z3} l_\beta + 2l_t l_x l_{z3} l_\beta + 2l_s^2 l_{z4} l_\beta + 2l_s l_x l_{z4} l_\beta - 2l_s l_{z3} l_{z4} l_\beta - \\
&2l_x l_{z3} l_{z4} l_\beta - 2l_s^2 l_\beta^2 - l_s l_t l_\beta^2 - 2l_s l_x l_\beta^2 - l_t l_x l_\beta^2 + l_s l_{z3} l_\beta^2 + l_x l_{z3} l_\beta^2 + l_s l_{z4} l_\beta^2 + \\
&l_x l_{z4} l_\beta^2 - \frac{2}{3} l_s l_\beta^3 + \frac{2}{3} l_T l_\beta^3 - \frac{5}{3} l_x l_\beta^3 - \frac{2}{3} l_{z3} l_\beta^3 + \Big(l_s^2 - \frac{19}{2} l_t^2 - \frac{3}{2} l_T^2 + 4l_T l_u - 4l_T l_x + \\
&5l_x^2 - 6l_T l_{z3} - 2l_x l_{z3} - 6l_{z3}^2 - 2l_T l_{z4} + 2l_x l_{z4} - 12l_{z3} l_{z4} - 7l_{z4}^2 + l_t(9l_T + 21l_x + \\
&16l_{z3} + 6l_{z4}) + 4l_T l_\beta - 4l_x l_\beta - 4l_{z3} l_\beta + l_s(-11l_t - 4l_T - 4l_x + 2l_{z3} + 10l_{z4} + \\
&2l_\beta) \Big) \zeta(2) + (7l_s - 8l_t + 4l_T - 3l_x + 2l_{z3} + 4l_{z4})\zeta(3) - \frac{41}{2}\zeta(4) + 4\mathrm{Li}_2\left(\frac{-u}{z_4}\right) \times \\
&l_T l_x - 2\mathrm{Li}_2(x) \Big(l_s^2 - l_x l_t + l_x l_{z3} - l_s l_t + 2l_s l_T - l_s l_{z3} \Big) + 2\mathrm{Li}_2^2(-x) + \\
&\mathrm{Li}_2\left(\frac{z_3}{z_4}\right) \Big(-l_s^2 + 2l_s(l_t - l_x) + 2l_x l_t + l_x^2 \Big) + \mathrm{Li}_2\left(\frac{m^2}{z_5}\right) \Big(-2l_s^2 - 4l_t l_T - 2l_x l_T + \\
&l_x^2 + 2l_s\,(2l_t - l_T - l_x) \Big) + \mathrm{Li}_2\left(\frac{T}{m^2}\right) \Big(-2\mathrm{Li}_2\left(-\frac{t}{s}\right) - 2\mathrm{Li}_2(-x) - \frac{11}{2} l_t^2 - 2l_T l_u \\
&-2l_x^2 - l_{z3}^2 - 2l_{z3} l_{z4} - l_{z4}^2 - l_t l_T + 6l_t(l_{z3} + l_{z4}) - 14\zeta(2) \Big) + \mathrm{Li}_2\left(\frac{m^2 s}{D}\right) \Big(3l_s^2 - \\
&l_t^2 - 2l_s(3l_t - l_T + l_u) + 2l_t(3l_T + l_u) + 2(l_T - l_x)l_x - 8\zeta(2) \Big) + \\
&\mathrm{Li}_2\left(-\frac{t}{s}\right) \Big(3l_s^2 - 2l_s(3l_t - 2l_T + l_u) - 2\Big(l_t^2 + 2l_T^2 - l_t(4l_T + l_u) + l_x^2 \Big) - 8\zeta(2) \Big) \\
&+\mathrm{Li}_2\left(-\frac{D}{m^2 t}\right) \Big(-2\mathrm{Li}_2\left(\frac{T}{m^2}\right) - 4l_s l_t - 2l_t^2 + 6l_t l_T - l_x^2 - 4\zeta(2) \Big) - \\
&\mathrm{Li}_2\left(\frac{T}{z_3}\right) \Big(2\mathrm{Li}_2(-x) + l_s^2 - 4l_s l_t + 2l_t(l_t + 3l_T - 2l_x - l_{z3} - 2l_{z4}) + 2\zeta(2) \Big) + \\
&\mathrm{Li}_2\left(\frac{m^2 x}{-T}\right) \Big(-l_x^2 + 2l_t(-l_t - 3l_T - 3l_x + l_{z3} + 2l_{z4}) - 2\zeta(2) \Big) + \\
&\mathrm{Li}_2\left(\frac{-t(1-\beta)}{2\,m^2}\right) \Big(-2\mathrm{Li}_2(-x) - 2l_s^2 - 4l_t l_T - 2l_x l_T + l_x^2 + 2l_s(2l_t - l_T + l_x) + \\
&8\zeta(2)) + \mathrm{Li}_2(-x) \Big(-\frac{9}{4} l_s^2 - 8l_x l_T + \frac{7}{4} l_x^2 - 2l_x l_{z3} + 9l_t l_x - 2l_t l_{z3} + 5l_s l_t - 6l_s l_T + \\
&\frac{7}{2} l_s l_x + 2l_s l_{z3} + 12\zeta(2) \Big) + \mathrm{Li}_3\left(\frac{m^2}{-t}\right) (-6l_s - l_t + l_T + 2l_x - 4l_{z4}) +
\end{aligned}
$$

$$\mathrm{Li}_3\left(\frac{T}{m^2}\right)(-4l_s+8l_t+3l_T+2l_x-6l_{z3}-10l_{z4})+2\mathrm{Li}_3\left(\frac{z_3}{t}\right)(l_t+l_T-4l_x-$$

$$l_{z3}-2l_{z4})+2\left(\mathrm{Li}_3\left(\frac{m^2(1-\beta)}{2\,z_5}\right)-\mathrm{Li}_3\left(\frac{-t(1-\beta)}{2\,z_5}\right)\right)(l_t+3l_T+3l_x-l_{z3}-$$

$$2l_{z4})+2\mathrm{Li}_3\left(\frac{z_4}{t}\right)(l_t+l_T+5l_x-l_{z3}-2l_{z4})+2\mathrm{Li}_3\left(-\frac{u}{t}\right)(3l_s-3l_t-2l_T)+$$

$$2\left(\mathrm{Li}_3\left(-\frac{m^2t}{D}\right)-\mathrm{Li}_3\left(\frac{t^2}{D}\right)\right)(2l_s+l_t-3l_T)+2\left(\mathrm{Li}_3\left(-\frac{m^2u}{sz_5}\right)-2\mathrm{Li}_3\left(\frac{z_3}{-u}\right)\right.$$

$$\left.+\mathrm{Li}_3\left(-\frac{sz_6}{m^2u}\right)-2\mathrm{Li}_3\left(\frac{-u}{z_4}\right)\right)(l_s-l_t-l_T)-2\mathrm{Li}_3\left(-\frac{D}{tu}\right)l_T-2\left(\mathrm{Li}_3\left(\frac{m^2}{z_5}\right)\right.$$

$$\left.+\mathrm{Li}_3\left(\frac{z_6}{m^2}\right)\right)(l_s-l_t+l_T)-2\mathrm{Li}_3\left(-\frac{t}{s}\right)(l_t-l_T)-2\mathrm{Li}_3\left(\frac{m^2s}{D}\right)(l_t-2l_T)+$$

$$2\mathrm{Li}_3(x)(3l_s-l_x)+2\mathrm{Li}_3\left(\frac{sxz_5}{m^2\,t}\right)(l_s-l_x)-2\mathrm{Li}_3\left(\frac{z_4(1-\beta)}{-2\,m^2}\right)(2l_s-l_t-l_T+$$

$$l_x)+2\mathrm{Li}_3\left(\frac{z_3}{z_4}\right)l_x+2\mathrm{Li}_3\left(\frac{z_6}{z_5}\right)l_s+2\left(\mathrm{Li}_3\left(\frac{t(1-\beta)}{2\,z_3}\right)+\mathrm{Li}_3\left(\frac{z_3}{s\beta}\right)+\mathrm{Li}_3\left(\frac{z_5}{t\beta}\right)\right)$$

$$\times(l_s+l_x)+2\mathrm{Li}_3\left(\frac{s(1-\beta)}{-2\,z_3}\right)(-2l_s+l_t+l_T+l_x)-2\mathrm{Li}_3\left(\frac{1-\beta}{-2\beta}\right)(l_s+4l_T-$$

$$5l_x-4l_{z3})+2\mathrm{Li}_3\left(\frac{-x^2}{1-x^2}\right)(l_s+2l_T-2l_x-2l_{z3})+6\mathrm{Li}_3\left(\frac{1-\beta}{2}\right)(l_T-l_x-$$

$$l_{z3})-2\mathrm{Li}_3\left(\frac{2\,z_6}{m^2(1+\beta)}\right)(2l_s-l_t-3l_T+2l_x+l_{z3}+2l_{z4})+2\mathrm{Li}_3\left(\frac{-2\,z_6}{t(1+\beta)}\right)$$

$$\times(2l_s-l_t-3l_T+2l_x+l_{z3}+2l_{z4})+2\mathrm{Li}_3(-x)(8l_s+4l_t-3l_T-7l_x-2l_{z3}+$$

$$4l_{z4})-3\mathrm{Li}_4\left(\frac{m^2}{-t}\right)-6\mathrm{Li}_4\left(-\frac{t}{s}\right)+3\mathrm{Li}_4\left(\frac{T}{m^2}\right)+16\mathrm{Li}_4\left(\frac{T}{t}\right)-12\mathrm{Li}_4\left(\frac{1-\beta}{2}\right)$$

$$-12\mathrm{Li}_4\left(\frac{1+\beta}{2}\right)+L_{-++}\left(1,0,0,\frac{m^2}{-T}\right)+\frac{9}{2}L_{-++}\left(1,0,0,-\frac{t}{T}\right)-$$

$$L_{-++}\left(1,0,0,\frac{-2}{1-\beta}\right)-L_{-++}\left(1,0,0,\frac{-2}{1+\beta}\right)-5L_{-++}\left(1,0,\frac{m^2}{-T},-1\right)-$$

$$2L_{-++}\left(1,0,\frac{m^2}{-T},\frac{1}{x}\right)-2L_{-++}\left(1,0,\frac{m^2}{-T},x\right)-\frac{3}{2}L_{-++}\left(1,\frac{m^2}{-T},\frac{m^2}{-T},0\right)-$$

$$4L_{-++}\left(1,\frac{m^2}{-T},\frac{m^2}{-T},\frac{u}{t}\right)+2L_{-++}\left(1,\frac{m^2}{-T},\frac{m^2}{-T},\frac{1}{x}\right)+2L_{-++}\left(1,\frac{m^2}{-T},\frac{m^2}{-T},x\right)$$

$$-3L_{-++}\left(\frac{t}{T},0,0,\frac{s}{t}\right)+2L_+\left(0,0,\frac{T}{m^2},\frac{1}{x}\right)+2L_+\left(0,0,\frac{T}{m^2},x\right)-$$

$$2L_+\left(0,0,\frac{T}{t},\frac{s}{t}\right)+2L_+\left(0,1,-1,\frac{1}{x}\right)+2L_+(0,1,-1,x)+4L_+(0,1,\frac{t}{s},\frac{-2}{1-\beta})$$

$$
\begin{aligned}
&+4L_{+}\left(0,1,\frac{t}{s},\frac{-2}{1+\beta}\right)+8L_{+}\left(0,1,\frac{1+\beta}{-2},-\frac{t}{T}\right)+8L_{+}\left(0,1,\frac{1-\beta}{-2},-\frac{t}{T}\right)+\\
&2L_{+}\left(0,\frac{m^2}{-t},\frac{T}{t},\frac{m^2}{-T}\right)-2L_{+}\left(0,\frac{m^2\,t}{-D},\frac{tT}{D},\frac{m^2}{-T}\right)+2L_{+}\left(0,\frac{m^2\,t}{-D},\frac{tT}{D},\frac{1}{x}\right)+\\
&2L_{+}\left(0,\frac{m^2\,t}{-D},\frac{t\,T}{D},x\right)+2L_{+}\left(0,-\frac{t}{s},\frac{t}{s},\frac{m^2}{-T}\right)-2L_{+}\left(0,\frac{t^2}{D},\frac{t\,T}{-D},-1\right)-\\
&4L_{+}\Bigg(0,\frac{T}{m^2},\frac{-T}{m^2},-1)+4L_{+}\left(0,\frac{T}{m^2},\frac{-T}{m^2},\frac{-2}{1-\beta}\right)+4L_{+}(0,\frac{T}{m^2},\frac{-T}{m^2},\frac{-2}{1+\beta}\Bigg)\\
&-6L_{+}\left(0,\frac{T}{t},-\frac{T}{t},-1\right)+4L_{+}\left(0,\frac{T}{t},-\frac{T}{t},\frac{1}{x}\right)+4L_{+}\left(0,\frac{T}{t},-\frac{T}{t},x\right)-\\
&2L_{+}\left(0,-\frac{u}{s},-\frac{t}{s},-1\right)+2L_{+}\left(0,-\frac{u}{s},-\frac{t}{s},\frac{1}{x}\right)+2L_{+}\left(0,-\frac{u}{s},-\frac{t}{s},x\right)-\\
&4L_{+}\left(0,\frac{T}{z_6},\frac{T(1+\beta)}{-2\,z_6},\frac{-2}{1-\beta}\right)-4L_{+}\left(0,\frac{T}{z_6},\frac{T(1+\beta)}{-2\,z_6},\frac{-2}{1+\beta}\right)+\\
&2L_{+}\left(0,\frac{1-\beta}{2},\frac{1+\beta}{2},\frac{1}{x}\right)+2L_{+}\left(0,\frac{1-\beta}{2},\frac{1+\beta}{2},x\right)+\\
&2L_{+}\left(0,\frac{m^2(1-\beta)}{2\,z_5},\frac{T(1-\beta)}{-2\,z_5},\frac{1}{x}\right)+2L_{+}\left(0,\frac{m^2(1-\beta)}{2\,z_5},\frac{T(1-\beta)}{-2\,z_5},x\right)+\\
&4L_{+}\left(0,\frac{T(1-\beta)}{2\,z_6},\frac{T(1+\beta)}{2\,z_6},-1\right)-2L_{+}\left(0,\frac{T(1-\beta)}{2\,z_6},\frac{T(1+\beta)}{2\,z_6},x\right)+\\
&2L_{+}\left(0,\frac{1+\beta}{2},\frac{1-\beta}{2},\frac{1}{x}\right)+2L_{+}\left(0,\frac{1+\beta}{2},\frac{1-\beta}{2},x\right)-2L_{+}\left(\frac{m^2}{-T},0,-x,x\right)\\
&-4L_{+}\left(\frac{m^2}{-T},1,-1,\frac{1}{x}\right)-2L_{+}\left(\frac{m^2}{-T},\frac{T}{t},-\frac{T}{t},-1\right)-2L_{+}\left(\frac{m^2}{-T},\frac{T}{t},-\frac{T}{t},\frac{u}{t}\right)-\\
&4L_{+}\left(\frac{m^2}{-T},\frac{1-\beta}{2},\frac{1+\beta}{2},\frac{u}{t}\right)-4L_{+}\left(\frac{m^2}{-T},\frac{1-\beta}{2},\frac{1+\beta}{2},\frac{1}{x}\right)-\\
&4L_{+}\left(\frac{m^2}{-T},\frac{1+\beta}{2},\frac{1-\beta}{2},\frac{u}{t}\right)-4L_{+}\left(\frac{m^2}{-T},\frac{1+\beta}{2},\frac{1-\beta}{2},\frac{1}{x}\right)-\\
&4L_{+}\left(\frac{1}{x},\frac{1-\beta}{2},\frac{1-\beta}{-2},-1\right)-4L_{+}\left(x,\frac{1+\beta}{2},\frac{1+\beta}{-2},-1\right)-\\
&L_{+++}\left(0,0,\frac{m^2}{-T},-1\right)+2L_{+++}\left(0,0,\frac{m^2}{-T},\frac{1}{x}\right)+2L_{+++}\left(0,0,\frac{m^2}{-T},x\right)-\\
&2L_{+++}\left(0,\frac{m^2}{-T},\frac{u}{t},-1\right)-2L_{+++}\left(0,\frac{m^2}{-T},\frac{u}{t},\frac{m^2}{-T}\right)+2L_{+++}\left(0,\frac{m^2}{-T},\frac{u}{t},\frac{1}{x}\right)+\\
&2L_{+++}\left(0,\frac{m^2}{-T},\frac{u}{t},x\right)+2L_{+++}\left(0,\frac{m^2}{-T},\frac{1}{x},\frac{1}{x}\right)+L_{+++}\left(0,\frac{1}{x},\frac{1}{x},\frac{m^2}{-T}\right)-
\end{aligned}
$$

$$L_{+++}\left(\frac{m^2}{-T},\frac{m^2}{-T},0,\frac{1}{x}\right) - L_{+++}\left(\frac{m^2}{-T},\frac{m^2}{-T},0,x\right) - L_{+++}\left(\frac{1}{x},\frac{1}{x},0,\frac{m^2}{-T}\right)\Bigg];$$

$$\begin{aligned}
\operatorname{Im} D_2^{(-1)} &= \pi/(st), \qquad\qquad \operatorname{Im} D_2^{(0)} = -2\pi l_t/(st), \qquad (4.68)\\
\operatorname{Im} D_2^{(1)} &= \frac{\pi}{st}\Bigg[-\frac{3}{4}l_s^2 + 2l_t^2 - \frac{3}{4}l_x^2 - l_x l_{z3} - l_t l_x - 2l_t l_{z3} + l_x l_{z4} - l_{z4}^2 + \\
&\quad l_s(l_t + \frac{l_x}{2} + l_{z3} + l_{z4}) - 2\zeta(2) - 2\mathrm{Li}_2\left(\frac{m^2}{z_5}\right) - 2\mathrm{Li}_2\left(\frac{t(1-\beta)}{-2\,m^2}\right)\Bigg],\\
\operatorname{Im} D_2^{(2)} &= \frac{\pi}{st}\Bigg[\frac{7}{12}l_s^3 - \frac{l_x^3}{12} - 2l_x l_t l_{z3} - l_x^2 l_{z3} + l_{z4}^2 - l_s\Big(2l_t^2 - \frac{5}{4}l_x^2 - 2l_t(l_x + l_{z3}) - \\
&\quad 2l_x(l_{z3} - l_{z4}) - l_{z4}^2\Big) - l_s^2\left(\frac{3}{4}l_x + l_{z3} + l_{z4}\right) + 4(l_t + l_x)\zeta(2) + 2\mathrm{Li}_2(x)\,l_x + \\
&\quad 2\mathrm{Li}_2\left(\frac{t(1-\beta)}{-2\,m^2}\right)(l_s - l_x) + 2\mathrm{Li}_2\left(\frac{z_3}{z_4}\right)l_x + 2\mathrm{Li}_2\left(\frac{m^2}{z_5}\right)(l_s + l_x) - 2\mathrm{Li}_3(x) - \\
&\quad 4\mathrm{Li}_3\left(\frac{z_3}{t}\right) + 2\mathrm{Li}_3\left(\frac{z_3}{z_4}\right) - 4\mathrm{Li}_3\left(\frac{z_4}{t}\right) - 2\mathrm{Li}_3\left(\frac{z_6}{z_5}\right)\Bigg].
\end{aligned}$$



FIG. 4. Massive box D_3 with two massive propagators.

The diagram corresponding to the third four-point function D_3 with two massive propagators is shown in Fig. 4 where we write

$$D_3 \equiv D(-p_2, p_4, -p_1, 0, 0, m, m)\,. \tag{4.69}$$

The kernel (4.40) for D_3 can be written as

$$K = ac\tilde{t} + bd\tilde{u} + (c+d)^2, \tag{4.70}$$

where we have introduced the positive-valued dimensionless variable

$$\tilde{u} \equiv -\frac{u}{m^2}, \qquad \tilde{u} \geq 1, \qquad \tilde{s} \geq \tilde{u}. \tag{4.71}$$

For the Feynman parameterization we choose $\{a,b,c,d\}$ as $\{x_1(1-x_2),\, 1-x_1,\, x_1x_2(1-x_3),\, x_1x_2x_3\}$, which gives the following integrand:

$$x_1^2x_2\left[x_1^2x_2^2 + \tilde{t}x_1^2x_2(1-x_2)(1-x_3) + \tilde{u}x_1(1-x_1)x_2x_3\right]^{-2-\varepsilon}.$$

The above expression never becomes negative. Therefore, the entire result for the box D_3 does not have an imaginary part. One can set $\delta = 0$ in the kernel from the very beginning.

That the box D_3 posesses no imaginary part can be seen in a less technical way by appealing to the Landau-Cutkosky cutting rules. The diagram corresponding to the box D_3 shown in Fig. 4 does not admit of any cuts such that the cut lines of the diagram are on their mass shell simultaneously.

As before we obtain two terms after the first integration over x_3. They are

$$I^{D_3}_{x_1x_2} = \frac{x_1^{-1-2\varepsilon} x_2^{-1-\varepsilon} \left[x_2 + \tilde{t}(1-x_2))\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{u}(1-x_1) - \tilde{t}x_1(1-x_2)]}, \tag{4.72}$$

$$II^{D_3}_{x_1x_2} = -\frac{x_1^{-\varepsilon} x_2^{-1-\varepsilon} \left[x_1x_2 + \tilde{u}(1-x_1)\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{u}(1-x_1) - \tilde{t}x_1(1-x_2)]}. \tag{4.73}$$

Note that the denominators of $I^{D_3}_{x_1x_2}$ and $II^{D_3}_{x_1x_2}$ change sign on the integration path, while the numerators stay positive (i.e. the relevant integrals have branch cuts). This can easily be seen by considering the numerators and denominators of the above integrands at two particular values of the variable x_1, for instance at $x_1 = 0$ and $x_1 = 1$. This means that although the whole D_3 box integral does not have an imaginary part, the two terms in (4.72) and (4.73) separately give rise to unphysical, spurious imaginary contributions. Of course, these are artefacts of having split the result into two terms. On the one hand, this somehow complicates things. On the other hand, the cancellation of imaginary contributions in the sum of the two terms (4.72) and (4.73) will serve as a good check for our final result. To control the imaginary contributions we do the following replacement in the denominators in (4.72) and (4.73):

$$\tilde{u}(1-x_1) - \tilde{t}x_1(1-x_2) \to \tilde{u}(1-x_1) - \tau x_1(1-x_2), \qquad \tau \equiv \tilde{t} - i\delta.$$

We start with the x_2-integration of the term $I^{D_3}_{x_1x_2}$ Eq. (4.72):

$$I^{D_3}_{x_2} = -\frac{\left[x_2\left(x_2 + \tilde{t}\,(1-x_2)\right)\right]^{-1-\varepsilon}\ {}_2F_1(1, -2\varepsilon, 1-2\varepsilon, \frac{\tilde{u}+\tau-x_2\tau}{\tilde{u}})}{2\tilde{u}\,(1+\varepsilon)\,\varepsilon}. \tag{4.74}$$

The above expression is singular at the lower integration limit $x_2 = 0$ due to the term $x_2^{-1-\varepsilon}$. To find a subtraction term, we follow exactly the procedures defined after Eq. (4.56). Our subtraction term reads

$$I^{D_3,s}_{x_2} = \frac{x_2^{-1-\varepsilon}\,\tilde{t}^{-1-\varepsilon}\,[-1 - 2\varepsilon\,\ln\left(-\frac{\tau}{\tilde{u}}\right) + 4\varepsilon^2\mathrm{Li}_2\left(\frac{\tilde{u}+\tau}{\tilde{u}}\right) + 8\varepsilon^3\,\mathrm{Li}_3\left(\frac{\tilde{u}+\tau}{\tilde{u}}\right) + 16\varepsilon^4\mathrm{Li}_4\left(\frac{\tilde{u}+\tau}{\tilde{u}}\right)]}{2\tilde{u}\,(1+\varepsilon)\,\varepsilon}. \tag{4.75}$$

The above subtraction term can easily be integrated over x_2 to obtain

$$I^{D_3,s} = \frac{\tilde{t}^{-1-\varepsilon}\,\left[1 + 2\varepsilon\,\ln\left(-\frac{\tau}{\tilde{u}}\right) - 4\varepsilon^2\mathrm{Li}_2\left(\frac{\tilde{u}+\tau}{\tilde{u}}\right) - 8\varepsilon^3\,\mathrm{Li}_3\left(\frac{\tilde{u}+\tau}{\tilde{u}}\right) - 16\varepsilon^4\mathrm{Li}_4\left(\frac{\tilde{u}+\tau}{\tilde{u}}\right)\right]}{2\,(1+\varepsilon)\,\varepsilon^2\,\tilde{u}} \tag{4.76}$$

which can readily be expanded in ε. As the difference $I_{x_2}^{D_3} - I_{x_2}^{D_3,s}$ does not contain any poles, we can expand the difference in a series in ε and perform the analytical integration over the last variable x_2.

To integrate the term (4.73), we split $II_{x_1x_2}^{D_3}$ into two contributions, i.e.

$$\begin{aligned} II_{x_1x_2}^{D_3} &= -\frac{x_2^{-1-\varepsilon}\left[x_1x_2 + \tilde{u}(1-x_1)\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{u}(1-x_1) - \tau x_1(1-x_2)]} \\ &\quad -\frac{(x_1^{-\varepsilon}-1)x_2^{-1-\varepsilon}\left[x_1x_2 + \tilde{u}(1-x_1)\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{u}(1-x_1) - \tau x_1(1-x_2)]}. \end{aligned} \tag{4.77}$$

Then we integrate the first term in (4.77) over x_1 to obtain two hypergeometric functions which are expanded up to ε^4. As was done previously, we then introduce a subtraction term similar to (4.75) which is integrated analytically. The finite difference of the original integral and the subtraction term is then ready to be integrated over the last integration variable.

For the second term in (4.77) we introduce a subtraction term before the last two integrations:

$$II_{x_1x_2}^{D_3,s} = -\frac{(x_1^{-\varepsilon}-1)x_2^{-1-\varepsilon}\left[\tilde{u}(1-x_1)\right]^{-1-\varepsilon}}{(1+\varepsilon)[\tilde{u}(1-x_1) - \tau x_1]}, \tag{4.78}$$

which is obtained from the second term in (4.77) by the substitution $x_2 = 0$ in all terms except for $x_2^{-1-\varepsilon}$. We first trivially integrate out x_2 in (4.78) and expand the resulting expression in a series of ε. Because of the factor $(x_1^{-\varepsilon}-1)$ this expansion starts at the order ε^0. Thus, the subtraction term is finite and ready for the last integration.

As the difference of the second term in (4.77) and its subtraction term (4.78) does not contain any poles, we expand it up to ε^2 and integrate over x_1. Finally, collecting all the relevant pieces, we perform the last integration. Our result for the four-point box diagram D_3 reads:

$$\mathrm{Re}\, D_3^{(-2)} = 1/(tu), \qquad \mathrm{Re}\, D_3^{(-1)} = -[l_t + l_u]/(tu), \tag{4.79}$$

$$\begin{aligned} \mathrm{Re}\, D_3^{(0)} &= 2\left[l_tl_u - 2\,\zeta(2)\right]/(tu), \\ \mathrm{Re}\, D_3^{(1)} &= \frac{1}{t\,u}\Bigg[\frac{l_D^3}{3} - l_D^2l_t - \frac{l_t^3}{3} + 2l_t^2l_u - 2l_tl_Tl_u - 4l_tl_u^2 + l_u^3 + l_s\left(l_u^2 - l_t^2\right) + l_D\left(l_t^2 + l_u^2\right) - \\ &\quad 2\,l_u^2l_U + l_s^2(l_t - l_u) + 2(l_D - 3l_s + l_t + 3l_u)\zeta(2) - 4\zeta(3) + 2\mathrm{Li}_2\left(-\frac{t}{s}\right)(l_t - l_u) + \\ &\quad 2\mathrm{Li}_2\left(\frac{m^4}{D}\right)l_u - 2\mathrm{Li}_2\left(\frac{T}{m^2}\right)l_u - 2\mathrm{Li}_2\left(\frac{U}{m^2}\right)l_u + 2\mathrm{Li}_2\left(-\frac{D}{m^2t}\right)(l_u - l_t) + \\ &\quad 2\mathrm{Li}_3\left(\frac{m^4}{D}\right) - 2\mathrm{Li}_3\left(\frac{m^2}{-t}\right) - 2\mathrm{Li}_3\left(-\frac{m^2t}{D}\right) - 2\mathrm{Li}_3\left(\frac{m^2}{-u}\right) - 2\mathrm{Li}_3\left(-\frac{m^2u}{D}\right) + \\ &\quad 2\mathrm{Li}_3\left(-\frac{u}{t}\right)\Bigg], \end{aligned}$$

$$
\begin{aligned}
\mathrm{Re}\,D_3^{(2)} \quad = \quad & \frac{1}{t\,u}\Bigg[\frac{l_D^4}{3} - l_D^2 l_s^2 - \frac{5}{6}\,l_s^4 - \frac{4}{3}\,l_D^3 l_t + 4 l_D l_s^2 l_t + \frac{1}{6}\,l_s^3 l_t + 4 l_D^2 l_t^2 - 5 l_D l_s l_t^2 - \frac{3}{4}\,l_s^2 l_t^2 - \\
& \frac{11}{3}\,l_D l_t^3 + \frac{3}{2}\,l_s l_t^3 + \frac{17}{6}\,l_t^4 + \frac{1}{3}\,l_D^3 l_T - 2 l_D l_s^2 l_T + l_s^3 l_T + l_D^2 l_t l_T + 4 l_D l_s l_t l_T - \\
& 2 l_s^2 l_t l_T - 4 l_D l_t^2 l_T + 3 l_s l_t^2 l_T + l_s^2 l_T^2 - 2 l_s l_t l_T^2 - \frac{4}{3}\,l_D^3 l_u + 2 l_D^2 l_s l_u - l_D l_s^2 l_u + \\
& \frac{5}{2}\,l_s^3 l_u - l_D^2 l_t l_u - 2 l_D l_s l_t l_u - 2 l_s^2 l_t l_u + 3 l_D l_t^2 l_u + l_s l_t^2 l_u - \frac{13}{6}\,l_t^3 l_u - l_D^2 l_T l_u + \\
& 2 l_D l_s l_T l_u - 2 l_s l_t l_T l_u + l_t^2 l_T l_u - l_s l_T^2 l_u - \frac{1}{2}\,l_D^2 l_u^2 + l_D l_s l_u^2 - \frac{9}{4}\,l_s^2 l_u^2 + 2 l_D l_t l_u^2 + \\
& 2 l_s l_t l_u^2 - \frac{7}{4}\,l_t^2 l_u^2 - l_D l_T l_u^2 - l_s l_T l_u^2 + 5 l_t l_T l_u^2 + \frac{1}{2}\,l_T^2 l_u^2 - \frac{4}{3}\,l_D l_u^3 + \frac{1}{6}\,l_s l_u^3 + \\
& \frac{3}{2}\,l_t l_u^3 - \frac{19}{12}\,l_u^4 + 2 l_D^2 l_u l_U - 2 l_D l_s l_u l_U + 2 l_s^2 l_u l_U - 2 l_D l_t l_u l_U + 2 l_t^2 l_u l_U + \\
& 2 l_D l_T l_u l_U - 4 l_t l_T l_u l_U - 2 l_s l_u^2 l_U + 2 l_t l_u^2 l_U + l_T l_u^2 l_U + \frac{13}{3}\,l_u^3 l_U + l_D l_u l_U^2 + \\
& l_s l_u l_U^2 - 2 l_t l_u l_U^2 - l_T l_u l_U^2 - 2 l_u^2 l_U^2 - \frac{2}{3}\,l_u l_U^3 + \Big(7 l_D^2 - 2 l_D (3 l_s + 4 l_t - l_T + 2 l_u) - \\
& \frac{17}{2}\,l_s^2 - \frac{3}{2}\,l_t^2 + 3 l_s (7 l_t + 6 l_u) - 2 l_t (2 l_T + 7 l_u) + l_T^2 - 2 l_T l_u - 7 l_u^2 - 2 l_u l_U\Big)\,\zeta(2) + \\
& (-3 l_s + 21\,l_t - 2 l_T - 8 l_u + 4 l_U)\,\zeta(3) - \frac{57}{4}\,\zeta(4) + \mathrm{Li}_2\left(\frac{U}{m^2}\right)\Big(l_D^2 + 2 l_s^2 - 2 l_D l_t + \\
& 3 l_t^2 - 2 l_t l_T + l_T^2 + 2 l_t l_u + 2 l_T l_u + 3 l_u^2 - 2 l_s (l_t + l_u) - 2 l_u l_U - 2\zeta(2)\Big) - \\
& \mathrm{Li}_2\left(\frac{m^4}{D}\right) l_u^2 + \mathrm{Li}_2\left(-\frac{t}{s}\right)\left(2\mathrm{Li}_2\left(\frac{T}{m^2}\right) + 2\mathrm{Li}_2\left(\frac{U}{m^2}\right) - 5 l_s^2 - \frac{3}{2}\,l_t^2 + 2 l_t l_T - \right. \\
& \left. 6 l_t l_u + \frac{5}{2}\,l_u^2 + 5 l_s (l_t + l_u) + 4\zeta(2)\right) + \mathrm{Li}_2\left(-\frac{D}{m^2 t}\right)\left(2\mathrm{Li}_2\left(\frac{T}{m^2}\right) + 2\mathrm{Li}_2\left(\frac{U}{m^2}\right) - \right. \\
& \left. 2 l_s^2 - 2 l_D l_t + 3 l_t^2 + 2 l_t l_T - 2 l_t l_u - 4 l_u^2 + 2 l_s (l_t + l_u) + 4 l_u l_U + 4\zeta(2)\right) + \\
& \mathrm{Li}_2\left(\frac{m^2 s}{D}\right)\left(2\mathrm{Li}_2\left(\frac{T}{m^2}\right) + l_t^2 - l_u^2 + 2 l_u l_U + 6\zeta(2)\right) + \mathrm{Li}_2\left(\frac{T}{m^2}\right)\left(\mathrm{Li}_2\left(\frac{T}{m^2}\right) + \right. \\
& \left. 4\mathrm{Li}_2\left(\frac{U}{m^2}\right) + 2 l_D^2 + l_s^2 + 3 l_t^2 - 2 l_D (l_s + 2 l_t) + 2 l_s l_T + l_u^2 + 2 l_u l_U + 6\zeta(2)\right) + \\
& 2\mathrm{Li}_3\left(\frac{m^4}{D}\right)(l_D - l_t + l_u) + 2\left(\mathrm{Li}_3\left(-\frac{m^2 t}{T u}\right) - \mathrm{Li}_3\left(\frac{m^2 s}{D}\right)\right)(l_s - l_t - l_u) - \\
& 2\mathrm{Li}_3\left(-\frac{m^2 t}{D}\right)(l_D - l_s + l_T + l_u) - 2\mathrm{Li}_3\left(\frac{U}{m^2}\right)(l_T - l_u) - \mathrm{Li}_3\left(-\frac{t}{s}\right)(6 l_s + l_t - \\
& 7 l_u) + 2\mathrm{Li}_3\left(\frac{m^2}{-u}\right)(l_D + l_s + 2 l_t - 7 l_T + 2 l_u - 2 l_U) - 2\mathrm{Li}_3\left(-\frac{m^2 u}{D}\right)(l_D - l_s +
\end{aligned}
$$

$$
\begin{aligned}
& l_T) - 2\left(\mathrm{Li}_3\left(\frac{t^2}{D}\right) + \mathrm{Li}_3\left(\frac{D}{u^2}\right)\right)(l_s - l_t - l_T) - \mathrm{Li}_3\left(-\frac{u}{t}\right)(3l_s + l_t + 2l_T + 2l_u) \\
& -2\mathrm{Li}_3\left(\frac{D}{sU}\right) l_u + 4\mathrm{Li}_3\left(-\frac{D}{tU}\right) l_u + 2\mathrm{Li}_3\left(\frac{m^4}{TU}\right) l_u - 2\mathrm{Li}_3\left(-\frac{t^2}{sT}\right)(l_s - l_t) + \\
& 2\mathrm{Li}_3\left(\frac{T}{m^2}\right)(2l_D + l_s - 3l_t - 2l_T) + 2\left(\mathrm{Li}_3\left(\frac{m^2}{-t}\right) + \mathrm{Li}_3\left(\frac{-D}{tu}\right)\right)(l_s - l_t - 2l_T) \\
& +2\mathrm{Li}_4\left(\frac{m^2}{-t}\right) + \mathrm{Li}_4\left(-\frac{t}{s}\right) - 2\mathrm{Li}_4\left(\frac{T}{m^2}\right) + 10\,\mathrm{Li}_4\left(\frac{T}{t}\right) - 8\mathrm{Li}_4\left(\frac{m^2}{-u}\right) + \\
& 7\mathrm{Li}_4\left(-\frac{u}{s}\right) - 4\mathrm{Li}_4\left(\frac{U}{m^2}\right) - 4\mathrm{Li}_4\left(\frac{U}{u}\right) + L_{-++}\left(1, \frac{m^2}{-T}, \frac{m^2}{-T}, \frac{-U}{m^2}\right) - \\
& 2L_{-++}\left(1, \frac{m^2}{-T}, \frac{u}{t}, 0\right) + 2L_{-++}\left(1, \frac{m^2}{-T}, \frac{u}{t}, \frac{m^2}{-T}\right) - \frac{1}{2}\,L_{-++}\left(1, \frac{t}{u}, \frac{t}{u}, 0\right) - \\
& 2L_{-++}\left(1, \frac{m^2}{-T}, \frac{u}{t}, \frac{-U}{m^2}\right) + \frac{1}{2}\,L_{-++}\left(1, \frac{u}{t}, \frac{u}{t}, 0\right) - \frac{5}{2}\,L_{-++}\left(-\frac{s}{t}, 0, 0, -1\right) + \\
& 3L_{-++}\left(\frac{t}{T}, 0, 0, -1\right) + 5L_{-++}\left(\frac{t}{T}, 0, 0, \frac{u}{m^2}\right) + \frac{1}{2}\,L_{-++}\left(-\frac{s}{u}, 0, 0, -1\right) - \\
& 2L_{+}\left(0, 0, -\frac{t}{s}, -1\right) + 2L_{+}\left(0, 0, \frac{T}{t}, -1\right) + 4L_{+}\left(0, 0, \frac{T}{t}, \frac{u}{m^2}\right) + \\
& 2L_{+}\left(0, 0, \frac{m^2}{-u}, \frac{u}{m^2}\right) - 2L_{+}\left(0, 1, \frac{t}{s}, -\frac{t}{T}\right) + 2L_{+}\left(0, 1, \frac{t}{s}, \frac{u}{m^2}\right) + \\
& 2L_{+}\left(0, \frac{t^2}{D}, \frac{tT}{-D}, -1\right) - 2L_{+}\left(0, \frac{t^2}{D}, \frac{tT}{-D}, -\frac{t}{T}\right) + 2L_{+}\left(0, \frac{t^2}{D}, \frac{tT}{-D}, \frac{u}{m^2}\right) + \\
& 2L_{+}\left(0, \frac{T}{m^2}, \frac{-T}{m^2}, \frac{s}{t}\right) - 2L_{+}\left(0, \frac{T}{m^2}, \frac{-T}{m^2}, -\frac{t}{T}\right) + 2L_{+}\left(\frac{m^2}{-T}, 0, \frac{T}{m^2}, \frac{u}{t}\right) - \\
& 2L_{+}\left(\frac{m^2}{-T}, \frac{U}{u}, \frac{m^2}{-u}, \frac{-U}{m^2}\right)\Bigg];
\end{aligned}
$$

$$
\mathrm{Im}\, D_3^{(j)} = 0. \tag{4.80}
$$

The non-planar topological structure of the four-point function D_3 implies that D_3 has to be $(t \leftrightarrow u)$–symmetric (see Fig. 4). This can best be seen by exchanging the momenta $p_3 \leftrightarrow p_4$ in Fig. 4 followed by a twist of the r.h.s. of Fig. 4 [3]. The $(t \leftrightarrow u)$–symmetry provides for a check on our results for D_3. The symmetry is obviously satisfied for $\mathrm{Re}D_3^{(-2)}$, $\mathrm{Re}D_3^{(-1)}$ and $\mathrm{Re}D_3^{(0)}$ but is not manifest for $\mathrm{Re}D_3^{(1)}$ and $\mathrm{Re}D_3^{(2)}$ in (4.79). However, it is quite straightforward to verify numerically that the $(t \leftrightarrow u)$–symmetry indeed holds for all coefficient functions in (4.79).

Apart from the internal checks mentioned earlier the most important check on our four-point function results has been a comparison with numerical results provided to us by M.M. Weber [17] for several phase space points. Within numerical errors we have found complete agreement with the results of M.M. Weber for each of the three four-point functions. It is important to emphasize that the approach of M.M. Weber to numerically evaluate the four-point functions is completely different from ours [18].

---

[3] The $(t \leftrightarrow u)$–symmetry is not so easy to see when exchanging $p_1 \leftrightarrow p_2$ in Fig. 4. In this case the $(t \leftrightarrow u)$–symmetry becomes apparent only after Feynman parametrization.

V. SUMMARY AND CONCLUSIONS

We have presented analytical results up to $\mathcal{O}(\varepsilon^2)$ for all the massive scalar one-loop integrals that arise in the calculation of one-loop matrix elements in heavy flavor hadroproduction. Many of our results are new (see Table 1). The one-loop scalar integrals are needed for that part of the NNLO hadroproduction of heavy flavours which is obtained from the product of one-loop contributions called loop-by-loop contribution.

What remains to be done in order to obtain the full one-loop amplitude structure is to take into account positive powers of ε (up to $\mathcal{O}(\varepsilon^2)$) resulting from the Passarino–Veltman decomposition and the spin algebra. The full one-loop amplitudes to order ε^0 were given in [13]. The missing results for the ε- and ε^2-coefficients of the one-loop amplitudes will be presented in a forthcoming publication [19]. In a last step the amplitudes themselves have to be squared, which, in the case of gluon-initiated production, will generate further positive powers of ε in dimensional regularization. The calculation of the loop-by-loop contributions in Fig. 1 is a necessary starting point in the evaluation of the NNLO contributions to heavy quark pair production in hadronic interactions. It is very likely that the calculation of the other three classes of diagrams in Fig. 1 will proof to be very difficult. This holds true in particular for the massive two-loop box contributions.

In the Laurent series expansion of the scalar one-loop integrals the successive coefficient functions increase in length and complexity with each order of ε. The reason is that the ε–expansion of the integrand before the last parametric integration itself generates coefficient functions with increasing complexity with each order of ε. The most complex expressions arise from the box contributions where one encounters multiple polylogarithms up to weight and depth four at $\mathcal{O}(\varepsilon^2)$.

In a numerical NNLO evaluation of heavy hadron production the various contributing pieces will have to be evaluated at many values of the kinematical variables. This requires efficiency in the numerical codes for each of the contributing pieces. We believe that we have provided for such numerical efficiency in the loop-by-loop portion of the NNLO calculation by presenting results in analytical form which are fast to evaluate numerically. All our results are available in electronic form [20]. We are planning to present our results in terms of multiple polylogarithms in the near future. In recent years number of new methods were developed for semi-numerical evaluation of general Feynman diagrams (see e.g. [18,21,22]). First numerical tests have shown that our representation in terms of the L-functions perform better than the present implementation of the flexible all-purpose approach described in [18,21].

The analytical results presented in this paper cover the whole kinematical domain with a single expression. They evaluate numerically very fast and efficiently. Further advantages of having the results in analytical form are that they allow one to investigate various limiting cases as well as their analyticity properties. Also, when analytical results are available the mathematical structure of the results becomes manifest which would not be visible in a purely numerical approach.

The full calculation of the NNLO corrections to heavy hadron production at hadron colliders will be a very difficult task to complete. It involves the calculation of very many Feynman diagrams of many different topologies. The problem is further complicated by the fact that heavy hadron production is a multi-scale problem with three mass scales provided by the kinematic variables s and t in the loop expressions, and the mass of the heavy quark. It is clear that an undertaking of this dimension will have to involve many theorists and cannot be done by a single group alone. In this sense the present calculation is a first step (or second step [23]) in the direction of obtaining NNLO results on heavy hadron production at hadron colliders. The present calculation allows one to obtain a first glimpse of the mathematical and computational complexity that is waiting for us in the full NNLO calculation. This complexity does in fact already reveal itself in terms of a very rich polylogarithmic and multiple polylogarithmic structure of the Laurent series expansion of the scalar one-loop integrals as shown in this paper.

ACKNOWLEDGMENTS

We are grateful to M.M. Weber for providing us with his numerical results on massive three– and four–point functions which served as another check on the correctness of our analytical results. We thank A. Davydychev, G. Heinrich and M. Kalmykov for communications, and G. Heinrich and J. Gegelia for discussions. Z. M. would like to thank the Particle Theory Group of the Institut für Physik, Universität Mainz for hospitality. The work of Z. M. was supported by a DFG (Germany) grant under contract 436 GEO 17/3/03. M. R. was supported by the DFG through the Graduiertenkolleg "Eichtheorien" at the University of Mainz.

APPENDIX A

In this Appendix we evaluate a special two-point integral which is needed for the calculation of the one-loop fermion self-energy diagram insertion into the massive external fermion line. This integral is also needed for the definitions of the fermion mass and wave function renormalization constants in the on-shell renormalization scheme. In particular, we need to evaluate the integral

$$I_1 \equiv B(p,0,m) = \mu^{2\varepsilon} \int \frac{d^nq}{(2\pi)^n} \frac{1}{q^2[(q+p)^2 - m^2]} \tag{A1}$$

up to $\mathcal{O}(p^2 - m^2)$. We therefore Taylor expand I_1 around $p^2 = m^2$:

$$\begin{aligned} I_1 &= I_1\Big|_{p^2=m^2} + \frac{dI_1}{dp^2}\Big|_{p^2=m^2}(p^2 - m^2) + \ldots \\ &\equiv E_0 + E_1(p^2 - m^2) + \ldots \end{aligned} \tag{A2}$$

Note that the expansion coefficients E_i in (A2) are functions of ε. The first coefficient E_0 is nothing but B_3 obtained in Section II. The second coefficient E_1 is proportional to the sum of a scalar and a vector integral obtained by differentiating I_1 w.r.t. p^μ. One obtains

$$E_1 = \frac{1}{2p_\mu}\frac{dI_1}{dp^\mu} = -I_2 - \frac{p^\mu I_{2\mu}}{m^2}, \tag{A3}$$

where

$$I_{\{2,2\mu\}} = \mu^{2\varepsilon}\int \frac{d^nq}{(2\pi)^n}\frac{\{1, q_\mu\}}{q^2[(q+p)^2 - m^2]^2} = iC_\varepsilon(m^2)\frac{1}{m^2}\left\{\frac{1}{2\varepsilon}, \frac{p_\mu}{1-2\varepsilon}\right\}. \tag{A4}$$

One finally has

$$I_1 = iC_\varepsilon(m^2)\frac{1}{\varepsilon(1-2\varepsilon)}\left(1 - \frac{p^2 - m^2}{2m^2}\right) + \mathcal{O}[(p^2 - m^2)^2]. \tag{A5}$$

The result for the integral I_1 in the form (A5) was used in [24] to evaluate external heavy quark self-energy diagrams and obtain heavy quark wave function renormalization constants in the NLO calculation.

APPENDIX B

In this Appendix we shall demonstrate how the L-functions introduced in Eqs. (3.20) and (3.21) are related to multiple polylogarithms as defined in [14]. Multiple polylogarithms are defined as a limit of Z-sums, e.g.

$$Li_{m_k,\ldots,m_1}(x_k, \ldots, x_1) = \lim_{n_1\to\infty} \sum_{n_1 > n_2 \ldots > n_k > 0} \frac{x_1^{n_1} x_2^{n_2} \ldots x_k^{n_k}}{n_1^{m_1} n_2^{m_2} \ldots n_k^{m_k}}. \tag{B1}$$

The number $w = m_1 + \ldots + m_k$ is called the weight and k is called the depth of the multiple polylogarithm. The power series (B1) is convergent for $|x_i| < 1$, and can be analytically continued via the iterated integral representation:

$$\begin{aligned} Li_{m_k,\ldots,m_1}(x_k, \ldots, x_1) = \int\limits_0^{x_1x_2\ldots x_k} &\left(\frac{dt}{t}\circ\right)^{m_1-1} \frac{dt}{x_2x_3\ldots x_k - t} \circ \\ &\left(\frac{dt}{t}\circ\right)^{m_2-1} \frac{dt}{x_3\ldots x_k - t} \circ \ldots \circ \left(\frac{dt}{t}\circ\right)^{m_k-1} \frac{dt}{1-t}, \end{aligned} \tag{B2}$$

where the following notation is used for the iterated integrals:

$$\int\limits_0^\lambda \frac{dt}{a_n - t} \circ ... \circ \frac{dt}{a_1 - t} = \int\limits_0^\lambda \frac{dt_n}{a_n - t_n} \int\limits_0^{t_n} \frac{dt_{n-1}}{a_{n-1} - t_{n-1}} \times ... \times \int\limits_0^{t_2} \frac{dt_1}{a_1 - t_1}. \tag{B3}$$

Note that the classical polylogarithms

$$\mathrm{Li}_n(z) \equiv \int\limits_0^z \frac{\mathrm{Li}_{n-1}(\xi)}{\xi} d\xi, \ n \geq 2; \qquad \mathrm{Li}_1(z) \equiv -\ln(1-z) \tag{B4}$$

are a subset of multiple polylogarithms. Examples of this statement can be found in the subsequent discussion.

We start by considering the single-index L-function Eq. (3.21):

$$L_{\sigma_1}(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = \int_0^1 dy \, \frac{\ln(\alpha_1 + \sigma_1 y)}{\alpha_4 + y} \mathrm{Li}_2(\alpha_2 + \alpha_3 y). \tag{B5}$$

After changing the integration variable $y = (t_1 - \alpha_2)/\alpha_3$ we get

$$L_{\sigma_1} = \int\limits_{\alpha_2}^{\alpha_2+\alpha_3} \frac{dt_1}{\alpha_3} \frac{\ln(\alpha_1 + \sigma_1 \frac{t_1 - \alpha_2}{\alpha_3})}{\alpha_4 + \frac{t_1 - \alpha_2}{\alpha_3}} \mathrm{Li}_2(t_1) = \int\limits_{\alpha_2}^{\alpha_2+\alpha_3} dt_1 \frac{\ln \frac{\sigma_1}{\alpha_3} + \ln(\frac{\alpha_1 \alpha_3}{\sigma_1} - \alpha_2 + t_1)}{\alpha_3 \alpha_4 - \alpha_2 + t_1} \mathrm{Li}_2(t_1). \tag{B6}$$

The integration interval can be split into two pieces, $[\alpha_2, 0]$ and $[0, \alpha_2 + \alpha_3]$. One can then write L_{σ_1} as a sum of four terms:

$$L_{\sigma_1} = \ln \frac{\sigma_1}{\alpha_3} \left(\int\limits_0^{\alpha_2+\alpha_3} - \int\limits_0^{\alpha_2} \right) \frac{dt_1}{\gamma + t_1} \mathrm{Li}_2(t_1) + \left(\int\limits_0^{\alpha_2+\alpha_3} - \int\limits_0^{\alpha_2} \right) dt_1 \frac{\ln(\alpha + t_1)}{\gamma + t_1} \mathrm{Li}_2(t_1), \tag{B7}$$

where we have introduced the notation

$$\alpha = \frac{\alpha_1 \alpha_3}{\sigma_1} - \alpha_2, \qquad \gamma = \alpha_3 \alpha_4 - \alpha_2. \tag{B8}$$

Regarding Eq. (B7) it is clear that there are only two different types of integrals to be dealt with:

$$\int\limits_0^{t_m} \frac{dt_1}{\gamma + t_1} \mathrm{Li}_2(t_1) \qquad \text{and} \qquad \int\limits_0^{t_m} dt_1 \frac{\ln(\alpha + t_1)}{\gamma + t_1} \mathrm{Li}_2(t_1). \tag{B9}$$

with upper limits $t_m = \alpha_2 + \alpha_3$ or $t_m = \alpha_2$. The first integral can be evaluated analytically in terms of standard logarithms and classical polylogarithms up to Li_3. However, the same integral can also be expressed in terms of multiple polylogarithms via the integral representation (B2), e.g.

$$\int\limits_0^{t_m} \frac{dt_1}{\gamma + t_1} \mathrm{Li}_2(t_1) = \int\limits_0^{t_m} \frac{dt_1}{\gamma + t_1} \int\limits_0^{t_1} \frac{dt_2}{t_2} \int\limits_0^{t_2} \frac{dt_3}{1 - t_3} = -Li_{2,1}\left(-\gamma, \frac{-t_m}{\gamma}\right). \tag{B10}$$

We now deal with the second integral in (B9). Consider the following multiple polylogarithm of weight four:

$$Li_{2,1,1}\left(-\gamma, \frac{\alpha}{\gamma}, \frac{t_m}{-\alpha}\right) = \int\limits_0^{t_m} \frac{dt_2}{-\alpha - t_2} \int\limits_0^{t_2} \frac{dt_1}{-\gamma - t_1} \mathrm{Li}_2(t_1) = \int\limits_0^{t_m} \frac{dt_1}{\gamma + t_1} \mathrm{Li}_2(t_1) \int\limits_{t_1}^{t_m} \frac{dt_2}{\alpha + t_2}$$

$$= \int\limits_0^{t_m} \frac{dt_1}{\gamma + t_1} \mathrm{Li}_2(t_1) \ln(\alpha + t_m) - \int\limits_0^{t_m} \frac{dt_1}{\gamma + t_1} \mathrm{Li}_2(t_1) \ln(\alpha + t_1). \quad \text{(B11)}$$

In the first step we have used the usual trick to change the order of integration. As already noted before (see Eq. (B10)) the first term on the second line can be expressed through a multiple polylogarithm of weight three. Thus one has

$$\int\limits_0^{t_m} dt_1 \frac{\ln(\alpha + t_1)}{\gamma + t_1} \mathrm{Li}_2(t_1) = -Li_{2,1,1}\left(-\gamma, \frac{\alpha}{\gamma}, \frac{t_m}{-\alpha}\right) - Li_{2,1}\left(-\gamma, \frac{-t_m}{\gamma}\right) \ln(\alpha + t_m). \quad \text{(B12)}$$

Finally, substituting Eqs. (B10) and (B12) into Eq. (B7) we arrive at the desired relation

$$\begin{aligned} L_{\sigma_1}(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = {} & Li_{2,1,1}\left(-\gamma, \frac{\alpha}{\gamma}, \frac{\alpha_2}{-\alpha}\right) - Li_{2,1,1}\left(-\gamma, \frac{\alpha}{\gamma}, \frac{\alpha_2 + \alpha_3}{-\alpha}\right) \\ & + Li_{2,1}\left(-\gamma, \frac{-\alpha_2}{\gamma}\right) \left(\ln \frac{\sigma_1}{\alpha_3} + \ln(\alpha + \alpha_2)\right) \\ & - Li_{2,1}\left(-\gamma, \frac{-\alpha_2 - \alpha_3}{\gamma}\right) \left(\ln \frac{\sigma_1}{\alpha_3} + \ln(\alpha + \alpha_2 + \alpha_3)\right), \end{aligned} \quad \text{(B13)}$$

where α and γ are defined in Eq. (B8).

Next we turn to the triple-index L-function Eq. (3.20)

$$L_{\sigma_1\sigma_2\sigma_3}(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = \int_0^1 dy \frac{\ln(\alpha_1 + \sigma_1 y) \ln(\alpha_2 + \sigma_2 y) \ln(\alpha_3 + \sigma_3 y)}{\alpha_4 + y} \quad \text{(B14)}$$

We again change the integration variable $y = (1 - \alpha_3 - t_1)/\sigma_3$ and obtain

$$L_{\sigma_1\sigma_2\sigma_3} = - \int\limits_{1-\alpha_3}^{1-\alpha_3-\sigma_3} \frac{dt_1}{\sigma_3} \ln(\gamma_1 - \frac{\sigma_1}{\sigma_3} t_1) \ln(\gamma_2 - \frac{\sigma_2}{\sigma_3} t_1) \frac{\ln(1 - t_1)}{\gamma_3 - t_1/\sigma_3}, \quad \text{(B15)}$$

where

$$\gamma_1 = \alpha_1 + \frac{\sigma_1}{\sigma_3}(1 - \alpha_3), \qquad \gamma_2 = \alpha_2 + \frac{\sigma_2}{\sigma_3}(1 - \alpha_3), \qquad \gamma_3 = \alpha_4 + \frac{1 - \alpha_3}{\sigma_3}. \quad \text{(B16)}$$

At this point we take $\sigma_1 = \sigma_2 = -\sigma_3 = 1$, which does not affect the generality of our further discussion. Splitting the integration interval in (B15) into two pieces as before, we arrive at

$$L_{++-} = \left(\int\limits_0^{1-\alpha_3-\sigma_3} - \int\limits_0^{1-\alpha_3} \right) dt_1 \ln(\gamma_1' + t_1) \ln(\gamma_2' + t_1) \frac{\ln(1 - t_1)}{\gamma_3' + t_1}, \quad \text{(B17)}$$

with

$$\gamma_1' = \alpha_1 - 1 + \alpha_3, \qquad \gamma_2' = \alpha_2 - 1 + \alpha_3, \qquad \gamma_3' = \alpha_4 - 1 + \alpha_3. \tag{B18}$$

Now consider the following multiple polylogarithm of weight four:

$$-Li_{1,1,1,1}\left(-\gamma_3, \frac{\gamma_2}{\gamma_3}, \frac{\gamma_1}{\gamma_2}, \frac{-t_m}{\gamma_1}\right) = \int_0^{t_m} \frac{dt_4}{\gamma_1 + t_4} \int_0^{t_4} \frac{dt_3}{\gamma_2 + t_3} \int_0^{t_3} \frac{dt_2}{\gamma_3 + t_2} \int_0^{t_2} \frac{dt_1}{1 - t_1} = \tag{B19}$$

$$-\int_0^{t_m} \frac{dt_4}{\gamma_1 + t_4} \int_0^{t_4} \frac{dt_3}{\gamma_2 + t_3} \int_0^{t_3} dt_2 \frac{\ln(1-t_2)}{\gamma_3 + t_2} = -\int_0^{t_m} \frac{dt_4}{\gamma_1 + t_4} \int_0^{t_4} dt_2 \frac{\ln(1-t_2)}{\gamma_3 + t_2} \int_{t_2}^{t_4} \frac{dt_3}{\gamma_2 + t_3} =$$

$$-\int_0^{t_m} dt_4 \frac{\ln(\gamma_2 + t_4)}{\gamma_1 + t_4} \int_0^{t_4} dt_2 \frac{\ln(1-t_2)}{\gamma_3 + t_2} + \int_0^{t_m} \frac{dt_4}{\gamma_1 + t_4} \int_0^{t_4} dt_2 \frac{\ln(1-t_2)\ln(\gamma_2 + t_2)}{\gamma_3 + t_2} =$$

$$-I'(t_m) + \int_0^{t_m} dt_2 \frac{\ln(1-t_2)\ln(\gamma_2 + t_2)}{\gamma_3 + t_2} \int_{t_2}^{t_m} \frac{dt_4}{\gamma_1 + t_4} =$$

$$-I'(t_m) + I''(t_m) - \int_0^{t_m} dt_2 \frac{\ln(\gamma_1 + t_2)\ln(\gamma_2 + t_2)\ln(1-t_2)}{\gamma_3 + t_2},$$

where we have introduced the notation

$$I'(t_m) = \int_0^{t_m} dt_4 \frac{\ln(\gamma_2 + t_4)}{\gamma_1 + t_4} \int_0^{t_4} dt_2 \frac{\ln(1-t_2)}{\gamma_3 + t_2}, \tag{B20}$$

$$I''(t_m) = \ln(\gamma_1 + t_m) \int_0^{t_m} dt_2 \frac{\ln(1-t_2)\ln(\gamma_2 + t_2)}{\gamma_3 + t_2}.$$

The third term on the last line of (B19) is the integral of the required type needed to express L_{++-} in Eq. (B17). We can write:

$$\int_0^{t_m} dt_1 \ln(\gamma_1 + t_1)\ln(\gamma_2 + t_1)\frac{\ln(1-t_1)}{\gamma_3 + t_1} = Li_{1,1,1,1}\left(-\gamma_3, \frac{\gamma_2}{\gamma_3}, \frac{\gamma_1}{\gamma_2}, \frac{-t_m}{\gamma_1}\right) - I'(t_m) + I''(t_m). \tag{B21}$$

To express the last two terms of (B21) in terms of multiple polylogarithms we proceed as follows. Consider first

$$-Li_{1,1,1}\left(-\gamma_3, \frac{\gamma_2}{\gamma_3}, \frac{-t_m}{\gamma_2}\right) = \int_0^{t_m} \frac{dt_2}{\gamma_2 + t_2} \int_0^{t_2} dt_1 \frac{\ln(1-t_1)}{\gamma_3 + t_1} = \int_0^{t_m} dt_1 \frac{\ln(1-t_1)}{\gamma_3 + t_1} \int_{t_1}^{t_m} \frac{dt_2}{\gamma_2 + t_2} = \tag{B22}$$

$$\ln(\gamma_2 + t_m) \int_0^{t_m} dt_1 \frac{\ln(1-t_1)}{\gamma_3 + t_1} - \int_0^{t_m} dt_1 \frac{\ln(1-t_1)\ln(\gamma_2 + t_1)}{\gamma_3 + t_1} =$$

$$\ln(\gamma_2 + t_m) Li_{1,1}\left(-\gamma_3, \frac{-t_m}{\gamma_3}\right) - \int_0^{t_m} dt_1 \frac{\ln(1-t_1)\ln(\gamma_2 + t_1)}{\gamma_3 + t_1},$$

from which we immediately conclude that

$$I''(t_m)/\ln(\gamma_1+t_m) = Li_{1,1,1}\left(-\gamma_3, \frac{\gamma_2}{\gamma_3}, \frac{-t_m}{\gamma_2}\right) + \ln\left(\gamma_2+t_m\right) Li_{1,1}\left(-\gamma_3, \frac{-t_m}{\gamma_3}\right). \tag{B23}$$

Note that the above multiple polylogarithms of weight three and weight two can also be expressed in terms of logarithms and classical polylogarithms by direct evaluation of the corresponding integrals.

For the first integral in (B20) we write

$$I'(t_m) = \int\limits_0^{t_m} dt_4 \frac{\ln(\gamma_2+t_4)}{\gamma_1+t_4} \int\limits_0^{t_4} dt_2 \frac{\ln(1-t_2)}{\gamma_3+t_2} = \int\limits_0^{t_m} dt_4 \frac{\ln(\gamma_2+t_4)}{\gamma_1+t_4} Li_{1,1}\left(-\gamma_3, \frac{-t_4}{\gamma_3}\right). \tag{B24}$$

On the other hand one has

$$\begin{aligned} Li_{1,1,1,1}\left(-\gamma_3, \frac{\gamma_1}{\gamma_3}, \frac{\gamma_2}{\gamma_1}, \frac{-t_m}{\gamma_2}\right) &= \int\limits_0^{t_m} \frac{dt_2}{\gamma_2+t_2} \int\limits_0^{t_2} \frac{dt_1}{\gamma_1+t_1} Li_{1,1}\left(-\gamma_3, \frac{-t_1}{\gamma_3}\right) = \\ & \int\limits_0^{t_m} \frac{dt_1}{\gamma_1+t_1} Li_{1,1}\left(-\gamma_3, \frac{-t_1}{\gamma_3}\right) \int\limits_{t_1}^{t_m} \frac{dt_2}{\gamma_2+t_2} = \\ \ln(\gamma_2+t_m) \int\limits_0^{t_m} \frac{dt_1}{\gamma_1+t_1} Li_{1,1}\left(-\gamma_3, \frac{-t_1}{\gamma_3}\right) &- \int\limits_0^{t_m} dt_1 \frac{\ln(\gamma_2+t_1)}{\gamma_1+t_1} Li_{1,1}\left(-\gamma_3, \frac{-t_1}{\gamma_3}\right) = \\ & - \ln(\gamma_2+t_m) Li_{1,1,1}\left(-\gamma_3, \frac{\gamma_1}{\gamma_3}, \frac{-t_m}{\gamma_1}\right) - I'(t_m). \end{aligned} \tag{B25}$$

We then conclude that

$$I'(t_m) = -Li_{1,1,1,1}\left(-\gamma_3, \frac{\gamma_1}{\gamma_3}, \frac{\gamma_2}{\gamma_1}, \frac{-t_m}{\gamma_2}\right) - \ln(\gamma_2+t_m) Li_{1,1,1}\left(-\gamma_3, \frac{\gamma_1}{\gamma_3}, \frac{-t_m}{\gamma_1}\right). \tag{B26}$$

Finally, substituting Eqs. (B23) and (B26) into Eq. (B21) we write down the result for the integral

$$\begin{aligned} \int\limits_0^{t_m} dt_1 \ln(\gamma_1+t_1)\ln(\gamma_2+t_1)\frac{\ln(1-t_1)}{\gamma_3+t_1} &= \ln(\gamma_1+t_m)\ln(\gamma_2+t_m) Li_{1,1}\left(-\gamma_3, \frac{-t_m}{\gamma_3}\right) + \\ \ln(\gamma_2+t_m) Li_{1,1,1}\left(-\gamma_3, \frac{\gamma_1}{\gamma_3}, \frac{-t_m}{\gamma_1}\right) &+ \ln(\gamma_1+t_m) Li_{1,1,1}\left(-\gamma_3, \frac{\gamma_2}{\gamma_3}, \frac{-t_m}{\gamma_2}\right) + \\ Li_{1,1,1,1}\left(-\gamma_3, \frac{\gamma_2}{\gamma_3}, \frac{\gamma_1}{\gamma_2}, \frac{-t_m}{\gamma_1}\right) &+ Li_{1,1,1,1}\left(-\gamma_3, \frac{\gamma_1}{\gamma_3}, \frac{\gamma_2}{\gamma_1}, \frac{-t_m}{\gamma_2}\right) \end{aligned} \tag{B27}$$

which demonstrates how the two integrals in Eq. (B17) representing L_{++-} are related to multiple polylogarithms. The discussion of the other cases for the $L_{\sigma_1\sigma_2\sigma_3}$ functions proceeds along similar lines.

APPENDIX C

In Appendix C we discuss properties and identities involving the single- and triple-index L-functions in Eqs. (3.20) and (3.21). There are two different categories of identities which we discuss in turn. We start by considering the simplest identities originating from symmetries related to permutations in the indices and arguments. We then present further identities based on integration-by-parts techniques.

1. Symmetry properties

We start with the single-index function $L_{\sigma_1}(\alpha_1,\alpha_2,\alpha_3,\alpha_4)$. One notices that a change of the integration variable $y \to 1-y$ results in the identity

$$L_{\sigma_1}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) = -L_{-\sigma_1}(\alpha_1+\sigma_1,\alpha_2+\alpha_3,-\alpha_3,-\alpha_4-1) \tag{C1}$$

which implies that L_- can always be related to L_+, and vice versa. We have thus written our results for the three-point and four-point functions in the main text only in terms of the L_+ functions.

Next we turn to the triple-index L-function. Note that $L_{\sigma_1\sigma_2\sigma_3}(\alpha_1,\alpha_2,\alpha_3,\alpha_4)$ is symmetric under permutations of any two pairs of indices and arguments $\{\sigma_i,\alpha_i\}$ and $\{\sigma_j,\alpha_j\}$ for $(i\neq j)$. The same change of variables as above $y\to 1-y$ results in

$$L_{\sigma_1\sigma_2\sigma_3}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) = -L_{-\sigma_1-\sigma_2-\sigma_3}(\alpha_1+\sigma_1,\alpha_2+\sigma_2,\alpha_3+\sigma_3,-\alpha_4-1). \tag{C2}$$

Therefore, from the eight functions L_{---}, L_{--+}, L_{-+-}, L_{+--}, L_{-++}, L_{+-+}, L_{++-}, L_{+++} only two are independent. We have chosen to write our results in terms of L_{-++} and L_{+++}.

2. Integration-by-parts identities

The triple- and single-index L-functions L_{+++}, L_{-++} and L_+ defined in Eqs. (3.20) and (3.21) have been devised such that they have neither branch cuts nor poles on the integration path $y\in[0,1]$. This also implies that the L_{+++}, L_{-++} and L_+ functions are real. Remember that the branch cuts for the ln and Li_2 functions are $(-\infty,0]$ and $(1,+\infty)$, respectively. The domains of the functions L_{+++}, L_{-++} and L_+ are

$$\begin{aligned}
L_{+++}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) \quad &: \quad \alpha_1>0,\alpha_2>0,\alpha_3>0,\alpha_4<-1 \quad \text{or} \quad \alpha_4>0;\\
L_{-++}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) \quad &: \quad \alpha_1>1,\alpha_2>0,\alpha_3>0,\alpha_4<-1 \quad \text{or} \quad \alpha_4>0;\\
L_{+}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) \quad &: \quad \alpha_1>0,\alpha_2\leq 1,\alpha_2+\alpha_3\leq 1,\alpha_3\neq 0,\alpha_4<-1 \quad \text{or} \quad \alpha_4>0.
\end{aligned} \tag{C3}$$

Looking at the definition of the triple-index L–function in (3.20) one concludes that the boundary points $\alpha_1=0$ and/or $\alpha_2=0$ and/or $\alpha_3=0$ can be included in the domain of definition for L_{+++}. The same holds true for $\alpha_1=1$ and/or $\alpha_2=0$ and/or $\alpha_3=0$ for L_{-++}. Also, from the definition of the single-index function L_+ in (3.21) on concludes that the boundary point $\alpha_1=0$ can be added to its domain of definition.

The points $\alpha_4 = \{-1, 0\}$ can also be included in the domain if the values taken by other parameters α_i guarantee the convergence of the integral. In what follows we assume everywhere in this appendix that the conditions (C3) are satisfied. Nevertheless, it is always possible to analytically continue the parameters to the complex plane.

In order to obtain integration-by-parts identities one makes use of the standard integration-by-parts formula

$$\int\limits_0^1 UV'dy = UV\Big|_0^1 - \int\limits_0^1 VU'dy\,. \tag{C4}$$

We start with the triple-index functions L_{-++} and L_{+++} defined in Eq. (3.20). Setting U equal to the numerator $[\ln(\alpha_1 + \sigma_1 y)\ln(\alpha_2 + \sigma_2 y)\ln(\alpha_3 + \sigma_3 y)]$ and V' equal to the remainder $(\alpha_4 + y)^{-1}$ we then arrive at

$$L_{+++}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) = \begin{cases} \alpha_4 > 0: & \ln(\alpha_1+y)\ln(\alpha_2+y)\ln(\alpha_3+y)\ln(\alpha_4+y)\Big|_0^1 \\ & -L_{+++}(\alpha_4,\alpha_2,\alpha_3,\alpha_1) - L_{+++}(\alpha_1,\alpha_4,\alpha_3,\alpha_2) \\ & -L_{+++}(\alpha_1,\alpha_2,\alpha_4,\alpha_3); \\ \alpha_4 < -1: & \ln(\alpha_1+y)\ln(\alpha_2+y)\ln(\alpha_3+y)\ln(-\alpha_4-y)\Big|_0^1 \\ & -L_{-++}(-\alpha_4,\alpha_2,\alpha_3,\alpha_1) - L_{-++}(-\alpha_4,\alpha_1,\alpha_3,\alpha_2) \\ & -L_{-++}(-\alpha_4,\alpha_1,\alpha_2,\alpha_3); \end{cases} \tag{C5}$$

and

$$L_{-++}(\alpha_1,\alpha_2,\alpha_3,\alpha_4) = \begin{cases} \alpha_4 > 0: & \ln(\alpha_1-y)\ln(\alpha_2+y)\ln(\alpha_3+y)\ln(\alpha_4+y)\Big|_0^1 \\ & -L_{+++}(\alpha_4,\alpha_2,\alpha_3,-\alpha_1) - L_{-++}(\alpha_1,\alpha_4,\alpha_3,\alpha_2) \\ & -L_{-++}(\alpha_1,\alpha_2,\alpha_4,\alpha_3); \\ \alpha_4 < -1: & \ln(\alpha_1-y)\ln(\alpha_2+y)\ln(\alpha_3+y)\ln(-\alpha_4-y)\Big|_0^1 \\ & -L_{-++}(-\alpha_4,\alpha_2,\alpha_3,-\alpha_1) \\ & +L_{-++}(\alpha_3+1,\alpha_1-1,-\alpha_4-1,-\alpha_2-1) \\ & +L_{-++}(\alpha_2+1,\alpha_1-1,-\alpha_4-1,-\alpha_3-1). \end{cases} \tag{C6}$$

For the second part of Eq.(C6) we have made use of relation (C2).

There are some special cases when some of the α_i take values on the boundary of the domain of definition where one can still make use of the identities (C5) and (C6) even if the conditions (C3) are not met. For example, for the case $\{\alpha_1 = 0, \alpha_4 = -1\}$ the identitity (C6) is still valid. There are similar special cases for further identities to be derived below.

The integration-by-parts identities for the single-index L_+ function are more involved. To prepare ourselves we first write down the derivative of the dilog function in the integrand of (3.21). One has

$$\frac{d\mathrm{Li}_2(\alpha_2 + \alpha_3 y)}{dy} = -\frac{\ln(1-\alpha_2-\alpha_3 y)}{\frac{\alpha_2}{\alpha_3} + y}. \tag{C7}$$

In the case of the single-index function it will prove important to consider two different choices for U. We start by setting the whole numerator $[\ln(\alpha_1+\sigma_1 y)\mathrm{Li}_2(\alpha_2+\alpha_3 y)]$ in the integrand of Eq. (3.21) to U. For V' we then have $(\alpha_4+y)^{-1}$. One obtains

$$L_+(\alpha_1,\alpha_2,\alpha_3,\alpha_4) =$$
$$\left\{\begin{array}{rl}
\alpha_4>0,\alpha_3>0: & \ln(\alpha_1+y)\mathrm{Li}_2(\alpha_2+\alpha_3 y)\ln(\alpha_4+y)\Big|_0^1 \\
& -L_+(\alpha_4,\alpha_2,\alpha_3,\alpha_1) \\
& +L_{-++}(\frac{1-\alpha_2}{\alpha 3},\alpha_1,\alpha_4,\frac{\alpha_2}{\alpha_3})+\ln(\alpha_3)\int\limits_0^1 \frac{\ln(\alpha_1+y)\ln(\alpha_4+y)}{\frac{\alpha_2}{\alpha_3}+y}dy; \\
\alpha_4>0,\alpha_3<0: & \ln(\alpha_1+y)\mathrm{Li}_2(\alpha_2+\alpha_3 y)\ln(\alpha_4+y)\Big|_0^1 \\
& -L_+(\alpha_4,\alpha_2,\alpha_3,\alpha_1) \\
& +L_{+++}(\frac{\alpha_2-1}{\alpha 3},\alpha_1,\alpha_4,\frac{\alpha_2}{\alpha_3})+\ln(-\alpha_3)\int\limits_0^1 \frac{\ln(\alpha_1+y)\ln(\alpha_4+y)}{\frac{\alpha_2}{\alpha_3}+y}dy; \\
\alpha_4<-1,\alpha_3>0: & \ln(\alpha_1+y)\mathrm{Li}_2(\alpha_2+\alpha_3 y)\ln(-\alpha_4-y)\Big|_0^1 \\
& +L_+(-\alpha_4-1,\alpha_2+\alpha_3,-\alpha_3,-\alpha_1-1) \\
& -L_{-++}(\alpha_1+1,\frac{1-\alpha_2-\alpha_3}{\alpha_3},-\alpha_4-1,-\frac{\alpha_2+\alpha_3}{\alpha_3}) \\
& +\ln(\alpha_3)\int\limits_0^1 \frac{\ln(\alpha_1+y)\ln(-\alpha_4-y)}{\frac{\alpha_2}{\alpha_3}+y}dy; \\
\alpha_4<-1,\alpha_3<0: & \ln(\alpha_1+y)\mathrm{Li}_2(\alpha_2+\alpha_3 y)\ln(-\alpha_4-y)\Big|_0^1 \\
& +L_+(-\alpha_4-1,\alpha_2+\alpha_3,-\alpha_3,-\alpha_1-1) \\
& +L_{-++}(-\alpha_4,\alpha_1,\frac{\alpha_2-1}{\alpha 3},\frac{\alpha_2}{\alpha_3})+\ln(-\alpha_3)\int\limits_0^1 \frac{\ln(\alpha_1+y)\ln(-\alpha_4-y)}{\frac{\alpha_2}{\alpha_3}+y}dy.
\end{array}\right. \tag{C8}$$

An additional condition for (C8) has to be explicated because it does not follow automatically from (C3), namely the parameters α_2 and α_3 are restricted by

$$\frac{\alpha_2}{\alpha_3}<-1 \quad \text{or} \quad \frac{\alpha_2}{\alpha_3}>0. \tag{C9}$$

The integrals in (C8) are simple enough to be evaluated in terms of classical polylogarithms up to Li_3. We do not provide explicit results for these integrations since they are rather lengthy and, in addition, depend on relations between the parameters.

A second choice for U in (C4) provides further identities for L_+. In this case one sets $\mathrm{Li}_2(\alpha_2+\alpha_3 y)$ to U and $\ln(\alpha_1+y)/(\alpha_4+y)$ to V'. To calculate V one has to differentiate between three cases for the set of parameters α_1 and α_4. One has

$$V=\int\frac{\ln(\alpha_1+y)}{\alpha_4+y}dy=\left\{\begin{array}{rl}
\alpha_1<\alpha_4: & \ln(\alpha_1+y)\ln(\frac{\alpha_4+y}{\alpha_4-\alpha_1})+\mathrm{Li}_2(\frac{\alpha_1+y}{\alpha_1-\alpha_4}); \\
\alpha_1>\alpha_4>0: & \ln(\alpha_1-\alpha_4)\ln(\alpha_4+y)-\mathrm{Li}_2(\frac{\alpha_4+y}{\alpha_4-\alpha_1}); \\
\alpha_4<-1: & \ln(\alpha_1-\alpha_4)\ln(-\alpha_4-y)-\mathrm{Li}_2(\frac{\alpha_4+y}{\alpha_4-\alpha_1}).
\end{array}\right. \tag{C10}$$

Using Eq. (C10) one then obtains

$$L_+(\alpha_1,\alpha_2,\alpha_3,\alpha_4) =$$

$$
\left\{
\begin{array}{rl}
\alpha_1 < \alpha_4, \alpha_3 > 0: & \mathrm{Li}_2(\alpha_2 + \alpha_3 y) \left(\ln(\alpha_1 + y) \ln(\tfrac{\alpha_4 + y}{\alpha_4 - \alpha_1}) + \mathrm{Li}_2(\tfrac{\alpha_1 + y}{\alpha_1 - \alpha_4})\right) \Big|_0^1 \\
& + L_{-++}(\tfrac{1-\alpha_2}{\alpha 3}, \alpha_1, \alpha_4, \tfrac{\alpha_2}{\alpha_3}) \\
& - L_{+}(\tfrac{1-\alpha_2-\alpha_3}{\alpha_3}, \tfrac{\alpha_1+1}{\alpha_1-\alpha_4}, \tfrac{1}{\alpha_4-\alpha_1}, -\tfrac{\alpha_2+\alpha_3}{\alpha_3}) + \\
& \int\limits_0^1 \frac{\ln(\alpha_3)\left[\mathrm{Li}_2(\frac{\alpha_1+y}{\alpha_1-\alpha_4}) + \ln(\alpha_1+y)\ln(\frac{\alpha_4+y}{\alpha_4-\alpha_1})\right] - \ln(\alpha_4-\alpha_1)\ln(\alpha_1+y)\ln(\frac{1-\alpha_2}{\alpha_3}-y)}{\frac{\alpha_2}{\alpha_3}+y} dy; \\
\alpha_1 < \alpha_4, \alpha_3 < 0: & \mathrm{Li}_2(\alpha_2 + \alpha_3 y) \left(\ln(\alpha_1 + y) \ln(\tfrac{\alpha_4 + y}{\alpha_4 - \alpha_1}) + \mathrm{Li}_2(\tfrac{\alpha_1 + y}{\alpha_1 - \alpha_4})\right) \Big|_0^1 \\
& + L_{+++}(\tfrac{\alpha_2-1}{\alpha 3}, \alpha_1, \alpha_4, \tfrac{\alpha_2}{\alpha_3}) \\
& + L_{+}(\tfrac{\alpha_2-1}{\alpha_3}, \tfrac{\alpha_1}{\alpha_1-\alpha_4}, \tfrac{1}{\alpha_1-\alpha_4}, \tfrac{\alpha_2}{\alpha_3}) + \\
& \int\limits_0^1 \frac{\ln(-\alpha_3)\left[\mathrm{Li}_2(\frac{\alpha_1+y}{\alpha_1-\alpha_4}) + \ln(\alpha_1+y)\ln(\frac{\alpha_4+y}{\alpha_4-\alpha_1})\right] - \ln(\alpha_4-\alpha_1)\ln(\alpha_1+y)\ln(\frac{\alpha_2-1}{\alpha_3}+y)}{\frac{\alpha_2}{\alpha_3}+y} dy; \\
\alpha_1 > \alpha_4 > 0, \alpha_3 > 0 \ : & \mathrm{Li}_2(\alpha_2 + \alpha_3 y) \left(\ln(\alpha_1 - \alpha_4) \ln(\alpha_4 + y) - \mathrm{Li}_2(\tfrac{\alpha_4 + y}{\alpha_4 - \alpha_1})\right) \Big|_0^1 \\
& + L_{+}(\tfrac{1-\alpha_2-\alpha_3}{\alpha_3}, \tfrac{\alpha_4+1}{\alpha_4-\alpha_1}, \tfrac{1}{\alpha_1-\alpha_4}, -\tfrac{\alpha_2+\alpha_3}{\alpha_3}) + \\
& \int\limits_0^1 \frac{-\ln(\alpha_3)\mathrm{Li}_2(\frac{\alpha_4+y}{\alpha_4-\alpha_1}) + \ln(\alpha_1-\alpha_4)\ln(\alpha_4+y)\ln(1-\alpha_2-\alpha_3 y)}{\frac{\alpha_2}{\alpha_3}+y} dy; \\
\alpha_4 < -1, \alpha_3 > 0 \ : & \mathrm{Li}_2(\alpha_2 + \alpha_3 y) \left(\ln(\alpha_1 - \alpha_4) \ln(-\alpha_4 - y) - \mathrm{Li}_2(\tfrac{\alpha_4 + y}{\alpha_4 - \alpha_1})\right) \Big|_0^1 \\
& + L_{+}(\tfrac{1-\alpha_2-\alpha_3}{\alpha_3}, \tfrac{\alpha_4+1}{\alpha_4-\alpha_1}, \tfrac{1}{\alpha_1-\alpha_4}, -\tfrac{\alpha_2+\alpha_3}{\alpha_3}) \\
& + \int\limits_0^1 \frac{-\ln(\alpha_3)\mathrm{Li}_2(\frac{\alpha_4+y}{\alpha_4-\alpha_1}) + \ln(\alpha_1-\alpha_4)\ln(-\alpha_4-y)\ln(1-\alpha_2-\alpha_3 y)}{\frac{\alpha_2}{\alpha_3}+y} dy; \\
\alpha_1 > \alpha_4 > 0, \alpha_3 < 0 \ : & \mathrm{Li}_2(\alpha_2 + \alpha_3 y) \left(\ln(\alpha_1 - \alpha_4) \ln(\alpha_4 + y) - \mathrm{Li}_2(\tfrac{\alpha_4 + y}{\alpha_4 - \alpha_1})\right) \Big|_0^1 \\
& - L_{+}(\tfrac{\alpha_2-1}{\alpha_3}, \tfrac{\alpha_4}{\alpha_4-\alpha_1}, \tfrac{1}{\alpha_4-\alpha_1}, \tfrac{\alpha_2}{\alpha_3}) + \\
& \int\limits_0^1 \frac{-\ln(-\alpha_3)\mathrm{Li}_2(\frac{\alpha_4+y}{\alpha_4-\alpha_1}) + \ln(\alpha_1-\alpha_4)\ln(\alpha_4+y)\ln(1-\alpha_2-\alpha_3 y)}{\frac{\alpha_2}{\alpha_3}+y} dy; \\
\alpha_4 < -1, \alpha_3 < 0 \ : & \mathrm{Li}_2(\alpha_2 + \alpha_3 y) \left(\ln(\alpha_1 - \alpha_4) \ln(-\alpha_4 - y) - \mathrm{Li}_2(\tfrac{\alpha_4 + y}{\alpha_4 - \alpha_1})\right) \Big|_0^1 \\
& - L_{+}(\tfrac{\alpha_2-1}{\alpha_3}, \tfrac{\alpha_4}{\alpha_4-\alpha_1}, \tfrac{1}{\alpha_4-\alpha_1}, \tfrac{\alpha_2}{\alpha_3}) + \\
& \int\limits_0^1 \frac{-\ln(-\alpha_3)\mathrm{Li}_2(\frac{\alpha_4+y}{\alpha_4-\alpha_1}) + \ln(\alpha_1-\alpha_4)\ln(-\alpha_4-y)\ln(1-\alpha_2-\alpha_3 y)}{\frac{\alpha_2}{\alpha_3}+y} dy.
\end{array}
\right.
\tag{C11}
$$

In deriving (C11) it is important to take into account condition (C9). As was the case in Eq. (C8) the integrals in (C11) can be evaluated in terms of classical polylogarithms up to Li_3.

There is also one special case of the last identity (C11) when the first and fourth arguments of the single-index L_+ function are equal, e.g. $\alpha_1 = \alpha_4$. In this case L_+ can be expressed only in terms of the functions L_{-++} or L_{+++} as follows:

$$L_+(\alpha_1, \alpha_2, \alpha_3, \alpha_1) =$$

$$\begin{cases} \alpha_3 > 0: & \frac{1}{2}\ln^2(\alpha_1+y)\mathrm{Li}_2(\alpha_2+\alpha_3 y)\Big|_0^1 + \frac{1}{2}L_{-++}(\frac{1-\alpha_2}{\alpha_3},\alpha_1,\alpha_1,\frac{\alpha_2}{\alpha_3}) \\ & +\frac{\ln(\alpha_3)}{2}\int\limits_0^1 \frac{\ln^2(\alpha_1+y)}{\frac{\alpha_2}{\alpha_3}+y}dy, \\ \alpha_3 < 0: & \frac{1}{2}\ln^2(\alpha_1+y)\mathrm{Li}_2(\alpha_2+\alpha_3 y)\Big|_0^1 + \frac{1}{2}L_{+++}(\frac{\alpha_2-1}{\alpha_3},\alpha_1,\alpha_1,\frac{\alpha_2}{\alpha_3})+ \\ & +\frac{\ln(-\alpha_3)}{2}\int\limits_0^1 \frac{\ln^2(\alpha_1+y)}{\frac{\alpha_2}{\alpha_3}+y}dy\,. \end{cases} \tag{C12}$$

The third and last identity for the L_+ function is obtained from the definition (3.21) without making direct use of the integration-by-parts identity Eq. (C4). Nevertheless it can still be called an integration-by-parts identity because it makes use of the well-known identity

$$\mathrm{Li}_2(z) = \zeta(2) - \ln(z)\ln(1-z) - \mathrm{Li}_2(1-z), \quad z \in C \tag{C13}$$

which in turn is derived from the definition of the Li_2–function (B4) with the help of the integration-by-parts identity (C4). After transforming $\mathrm{Li}_2(\alpha_2+\alpha_3 y)$ according to (C13) one gets

$$L_+(\alpha_1,\alpha_2,\alpha_3,\alpha_4) =$$

$$\begin{cases} \begin{matrix}\alpha_3 > 0 \\ 0 \le \alpha_2 \le 1, \alpha_2+\alpha_3 \le 1\end{matrix} & : \; -L_+(\alpha_1,1-\alpha_2,-\alpha_2,\alpha_4) - L_{-++}(\frac{1-\alpha_2}{\alpha_3},\alpha_1,\frac{\alpha_2}{\alpha_3},\alpha_4)+ \\ & \int\limits_0^1 \frac{\ln(\alpha_1+y)}{\alpha_4+y}\Big(\zeta(2) - \ln(\alpha_3)[\ln(\alpha_2+\alpha_3 y) + \ln(\frac{1-\alpha_2}{c} - y)]\Big)\,dy; \\ \begin{matrix}\alpha_3 < 0 \\ 0 \le \alpha_2 \le 1, 0 \le \alpha_2+\alpha_3\end{matrix} & : \; -L_+(\alpha_1,1-\alpha_2,-\alpha_2,\alpha_4) - L_{-++}(-\frac{\alpha_2}{\alpha_3},\alpha_1,\frac{\alpha_2-1}{\alpha_3},\alpha_4)+ \\ & \int\limits_0^1 \frac{\ln(\alpha_1+y)}{\alpha_4+y}\Big(\zeta(2) - \ln(-\alpha_3)[\ln(\alpha_2+\alpha_3 y) + \ln(\frac{\alpha_2-1}{c} + y)]\Big)\,dy. \end{cases} \tag{C14}$$

A few final comments are appropiate. In spite of the rather complicated appearance of the identities (C5), (C6), (C8), (C11), (C14) these turn out to be very useful to reduce the length of the results presented in the main text. The first step in the chain of reductions is to write everything in terms of the functions L_+, L_{-++} and L_{+++}. In a second step one uses the identities written down in this Appendix to find the set of arguments of L-functions for which the number of the functions L_+, L_{-++} and L_{+++} is minimal. We have devised several programs for the MATHEMATICA computer algebra system which automatically find minimal sets of the single- and triple-index L-functions. With the help of these programs we have been able to greatly reduce the number of L-functions appearing in our results and have thereby greatly reduced their length. The same was done for the logarithms and classical polylogarithms using standard one- and two-variable identities for classical polylogarithms as given e.g. in [25].

REFERENCES

[1] P. Nason, S. Dawson and R. K. Ellis, Nucl. Phys. **B303**, 607 (1988); **B327**, 49 (1989); **B335**, 260(E) (1990).

[2] W. Beenakker, H. Kuijf, W. L. van Neerven and J. Smith, Phys. Rev. D **40**, 54 (1989); W. Beenakker, W. L. van Neerven, R. Meng, G. A. Schuler, and J. Smith, Nucl. Phys. **B351**, 507 (1991).

[3] M. Glück, J.F. Owens and E. Reya, Phys. Rev. D **17**, 2324 (1978);
B. L. Combridge, Nucl. Phys. **B151** (1979) 429;
J. Babcock, D. Sivers and S. Wolfram, Phys. Rev. D **18**, 162 (1978);
K. Hagiwara and T. Yoshino, Phys. Lett. **80B**, 282 (1979);
L. M. Jones and H. Wyld, Phys. Rev. D **17**, 782 (1978);
H. Georgi *et al.*, Ann. Phys. (N.Y.) **114**, 273 (1978).

[4] F. Abe *et al.*, [CDF Collaboration], Phys. Rev. Lett. **71**, 500 (1993);
D. Acosta *et al.* [CDF Collaboration], Phys. Rev. D **65**, 052005 (2002) [ArXiv: hep-ph/0111359].

[5] S. Abachi *et al.* [D0 Collaboration], Phys. Rev. Lett. **74**, 3548 (1995).

[6] M. Cacciari, S. Frixione, M. L. Mangano, P. Nason, G. Ridolfi, JHEP **0407**, 033 (2004) [ArXiv: hep-ph/0312132].

[7] G. 't Hooft and M. Veltman, Nucl. Phys. **B44**, 189 (1972);
C. G. Bollini and J. J. Giambiagi, Phys. Lett. **40B**, 566 (1972);
J. F. Ashmore, Nuovo Cimento Lett. **4**, 289 (1972).

[8] K. G. Chetyrkin and F. V. Tkachov, Nucl. Phys. **B192**, 159 (1981);
F. V. Tkachov, Phys. Lett. **100B**, 65 (1981).

[9] E. E. Boos, A. I. Davydychev, Theor. Math. Phys. **89**, 1052 (1991);
A. I. Davydychev, J. Math. Phys. **33**, 358 (1992).

[10] A. I. Davydychev and M. Yu. Kalmykov, Nucl. Phys. **B605**, 266 (2001) [ArXiv: hep-th/0012189].

[11] A. I. Davydychev and M. Yu. Kalmykov, Nucl. Phys. **B699**, 3 (2004) [ArXiv: hep-th/0303162].

[12] J. Fleischer, F. Jegerlehner and O.V. Tarasov, Nucl. Phys. **B672**, 303 (2003) [ArXiv: hep-ph/0307113].

[13] J. G. Körner and Z. Merebashvili, Phys. Rev. D **66**, 054023 (2002) [ArXiv: hep-ph/0207054].

[14] A. B. Goncharov, Math. Res. Lett. **5**, 497 (1998) [available at http://www.math.uiuc.edu/K-theory/0297];
S. Moch, P. Uwer and S. Weinzierl, J. Math. Phys. **43**, 3363 (2002) [ArXiv: hep-ph/0110083].

[15] J. Vollinga and S. Weinzierl, hep-ph/0410259.

[16] U. Nierste, D. Müller, M. Böhm, Z. Phys. **C57**, 605 (1993).

[17] M.M. Weber, private communication.

[18] M.M. Weber, Acta Phys. Polon. **B35**, 2655 (2004) [ArXiv: hep-ph/0410166].

[19] J. G. Körner, Z. Merebashvili and M. Rogal. To be published.

[20] See EPAPS Document No. E-PRVDAQ-71-025507 for our analytical results for the integrals in MATHEMATICA format. A direct link to this document may be found in the online article's HTML reference section. The document may also be reached via the EPAPS homepage (http://www.aip.org/pubservs/epaps.html) or from ftp.aip.org in the directory /epaps/. See the EPAPS homepage for more information.

[21] A. Ferroglia, M. Passera, G. Passarino and S. Uccirati, Nucl. Phys. **B680**, 199 (2004) [ArXiv: hep-ph/0311186]; **B650**, 162 (2003) [ArXiv: hep-ph/0209219].

[22] T. Binoth and G. Heinrich, Nucl. Phys. **B680**, 375 (2004) [ArXiv: hep-ph/0305234].

[23] W. Bernreuther *et al.*, Nucl. Phys. **B706**, 245 (2005) [ArXiv: hep-ph/0406046].

[24] B. Kamal, Z. Merebashvili, and A.P. Contogouris, Phys. Rev. D **51**, 4808 (1995); **55**, 3229(E) (1997).

[25] L. Lewin, *Polylogarithms and Associated Functions* (North Holland, Amsterdam, 1981).